

The analytic continuation of hyperbolic space

Yunhi Cho and Hyuk Kim*

Abstract

We define and study an extended hyperbolic space which contains the hyperbolic space and de Sitter space as subspaces and which is obtained as an analytic continuation of the hyperbolic space. The construction of the extended space gives rise to a complex valued geometry consistent with both the hyperbolic and de Sitter space. Such a construction shed a light and inspires a new insight for the study of the hyperbolic geometry and Lorentzian geometry. We discuss the advantages of this new geometric model as well as some of its applications.

1 Introduction

The hyperbolic space is an independent geometric entity with an infinite diameter and infinite volume which is already complete in its own right. But if we look at the hyperbolic space as a unit disk in the Kleinian model, then using the same metric formula we have a Lorentzian space with constant curvature outside the unit disk. Furthermore we can even draw a geometric figure lying across the ideal boundary. We naturally expect on this space the generalization of the basic geometric notions such as angle, length, volumes, ..., etc, and the similar relation between them to those on the hyperbolic space. But we immediately have difficulties in defining and deriving those due to the Lorentzian nature of the metric and multi-valuedness of the analytic functions representing various geometric formulas.

In this paper we show there is a natural way of extending the geometry of hyperbolic space to Lorentzian part and set a foundation of a geometry which connects and unifies these two different geometries by an analytic continuation method on the Kleinian model. We call such a unified space an extended hyperbolic space since we start from the hyperbolic space and then continue analytically to the Lorentzian part. The purpose of studying an extended hyperbolic space is not only to give a proper geometry on the Lorentzian space as a continuation of hyperbolic geometry but also to give a new insight to the hyperbolic geometry itself by studying a geometric object lying across the ideal boundary - the proper study of such object would be impossible otherwise.

Some of the basic notions such as angle, length and geodesic triangle on the extended hyperbolic plane has been considered and studied so far through cross ratio (see [16]) and a

*The second author was supported by grant no.(R01-1999-000-00002-0) from the Basic Research Program of the Korea Science & Engineering Foundation

rather ad hoc combinatorial method (see [6]). In this paper we study the extended hyperbolic space in a more systematic way and discuss some of its applications. In Section 2, we set up the geometry as an analytic continuation going over the singularity of the hyperbolic metric at the ideal boundary, we view the hyperbolic metric as a limit of complex perturbation called an ϵ -metric which is a complex regular metric. And we define and study distance, angle, length and k -dimensional volumes on the extended space in Section 4. Then such geometric quantities are given rather naturally with complex numbers - they of course coincide with the usual real values for the quantities inside the hyperbolic space. If we consider a nice region lying across the ideal boundary, then the volume of the region is of a finite complex value while the volume of the hyperbolic part and the Lorentzian part of the region are both infinite. In fact the measure defined on the extended hyperbolic space which extends the usual hyperbolic volume (and Lorentz volume) is a finitely (but not countably) additive complex measure and we study some of its delicate and strange properties in Section 3.

We would like to describe some of the advantages and expectations of using the extended hyperbolic model. When we study hyperbolic geometry we naturally want to extend the geometric objects and notions over and beyond the ideal boundary. But when we try to compute geometric quantities, we come across with a confusion of choosing an appropriate value among infinite possibilities of multi-values even for a distance or an angle. The extended model provide us a fulfilling consistency without such confusion once we choose an analytic continuation which follows naturally after fixing an ϵ -approximation of the metric. For instance the understanding of 1-dimensional extended model quickly leads us to be able to define an angle as a complex numbers on the general semi-Riemannian manifolds. Generalizing an earlier work of Kellerhals[8], Ushijima studied the volume of a hyperbolic polyhedron obtained by truncating with the dual plane of an ultra ideal vertex of a tetrahedron which is lying across the ideal boundary. He showed in [19] that the volume of the truncated tetrahedron is the real part of the value obtained after formal application of the known volume formula of a hyperbolic tetrahedron to this tetrahedron. But for the imaginary part it is multi-valued and there remains a problem of choice and interpretation of its geometric meaning. The extended model determines the unique value of the imaginary part and explains its geometric meaning as the volume of the truncated portion or as the area of the truncated face. See Example 5.11.

As we work more with the extended model, we found that the model is a natural and fundamental geometric setting as it works beautifully in every aspect. The n -dimensional extended hyperbolic space $\mathbb{S}_H^n$ is simply the standard sphere $\mathbb{S}^n$ topologically but with the geometry coming from the unit sphere of the Minkowski space. Its geometry has many resemblance with the spherical geometry and we can obtain various results through this conceptual but concrete analogy. For instance we can derive the Gauss-Bonnet formula on the extended space (this was first shown in [2]) combinatorially using Euler method as we did for the 2-sphere $\mathbb{S}^2$ without computing integrals and then of course can be generalized to the higher dimensions. (See Proposition 5.4.) Furthermore we can extend the hyperbolic trigonometry to the extended space, which of course implies that we have the same trigonometry on the de Sitter space (the Lorentzian part) and even for an object sitting across the ideal boundary. In principle we can apply the same method used for the standard sphere to obtain the

corresponding results on the extended hyperbolic space, which in turn give rise to the results for the hyperbolic space as well as de Sitter space. It would be interesting to observe that the volume of $\mathbb{S}_H^n$ differs from that of $\mathbb{S}^n$ by i^n . (See Theorem 2.3.)

As another illustration, let us consider Santaló's formula [15] giving the relation between the volume of a simplex and its dual in $\mathbb{S}^3$ or $\mathbb{H}^3$, Milnor's relation [10] between a convex polyhedron P and its dual P^* in $\mathbb{S}^3$, and Suárez-Peiró's result [17] for a simplex and its dual in $\mathbb{H}^n$. Then we see that there is a slight discrepancy between $\mathbb{S}^3$ and $\mathbb{H}^3$:

$$\text{vol}(P) + \text{vol}(P^*) + \frac{1}{2} \sum a_i(\pi - A_i) = \pi^2 \quad (\mathbb{S}^3)$$

$$\text{vol}(P) + \text{vol}(P^*) - \frac{1}{2} \sum a_i(\pi - A_i) = 0 \quad (\mathbb{H}^3)$$

where a_i and A_i are edge lengths and dihedral angles of P . Here if we use the extended model $\mathbb{S}_H^3$ instead of $\mathbb{H}^3$, the proof is essentially identical with the case for $\mathbb{S}^3$ and the formula can be written for both cases in a unified way as follows.

$$\text{vol}(P) - \text{vol}(\text{II}) + \text{vol}(\text{III}) - \text{vol}(\text{IV}) + \text{vol}(P^*) = 0$$

For this expression, the volume of simplices of type III lying across the boundary play the crucial role which is also interpreted as mean curvature integral in $(\mathbb{S}^3)$ and $(\mathbb{H}^3)$. And Suárez-Peiró's result is then also expected to be generalized for both cases in a single identity as

$$\text{vol}(P) - \text{vol}(\text{II}) + \text{vol}(\text{III}) - \cdots + (-1)^{n+1} \text{vol}(P^*) = 0$$

Conceptually things are getting tremendously easier giving us inspirations and this is one of the real merits using the extended model.

Lastly here are some speculations and the directions for possible further developments and applications. If we use the extended model, all the geometric quantities become complex valued and we can ask whether this is related to complex invariants of hyperbolic manifolds such as volume and Chern-Simon invariant pair in dimension 3. Using the extended model we can derive all the trigonometry and precise elementary geometric formulas for the Lorentzian spherical space (or de Sitter space) and this will be useful to study the discrete group actions on the Lorentzian space. Since the hyperbolic space and the Lorentzian space are dual each other, it is obviously advantageous to study the both subjects simultaneously in a unified geometric setting. Similar constructions for other semi-Riemannian cases, complex hyperbolic and quaternionic hyperbolic cases would also be very interesting.

Acknowledgement The authors would like to thank to Hyeonbae Kang, Chong Kyu Han, Dohan Kim for help in the analysis. In particular, the idea of ϵ -approximation came across with the suggestion of Professor Kang when we have a discussion with him and then we formulate and develop as given in this paper.

2 Hyperbolic sphere containing hyperbolic space

Let $\mathbb{R}^{n,1}$ denote the real vector space $\mathbb{R}^{n+1}$ equipped with the bilinear form of signature $(n, 1)$,

$$\langle x, y \rangle = -x_0 y_0 + x_1 y_1 + \cdots + x_n y_n,$$

for all $x = (x_0, x_1, \dots, x_n)$, $y = (y_0, y_1, \dots, y_n)$. Then the hyperbolic spaces H_+^n and H_-^n , pseudo-sphere S_1^n and light cone L^n are defined by

$$\begin{aligned} H_+^n &:= \{x \in \mathbb{R}^{n,1} | \langle x, x \rangle = -1, \quad x_0 > 0\}, \\ H_-^n &:= \{x \in \mathbb{R}^{n,1} | \langle x, x \rangle = -1, \quad x_0 < 0\}, \\ S_1^n &:= \{x \in \mathbb{R}^{n,1} | \langle x, x \rangle = 1\}, \\ L^n &:= \{x \in \mathbb{R}^{n,1} | \langle x, x \rangle = 0\}. \end{aligned}$$

We already know that $H_\pm^n$ is a Riemannian manifold which has a constant sectional curvature -1 , and that S_1^n is a Lorentzian manifold (or semi-Riemannian of signature $(-, +, \dots, +)$) which has a constant sectional curvature 1 (see, for example, [13]). Now we consider the Kleinian projective model. By the radial projection π_1 with respect to the origin from H_+^n into $\{1\} \times \mathbb{R}^n$, we obtain the induced Riemannian metric on the ball in $\{1\} \times \mathbb{R}^n$ as follows ([1],[14]),

$$ds_K^2 = \left(\frac{\sum x_i dx_i}{1 - |x|^2} \right)^2 + \frac{\sum dx_i^2}{1 - |x|^2}.$$

We will denote the unit ball with this metric by $\mathbb{H}^n$. If we extend this metric beyond the unit ball using the same formula, this metric induces a semi-Riemannian structure on the outside of the unit ball in $\{1\} \times \mathbb{R}^n$ and this will be denoted by $\mathbb{L}^n$. In fact, if we compare this metric with the one induced from the Lorentzian space $S_1^n \cap \{x = (x_0, x_1, \dots, x_n) | x_0 > 0\}$, by the radial projection into $\{1\} \times \mathbb{R}^n$, then they differ only by sign -1 . This sign change of metric implies the sign change of the sectional curvature from $+1$ to -1 , which, of course, the curvature of the metric ds_K^2 . In this way, we obtain an *extended Kleinian model* with metric ds_K^2 defined on $\{1\} \times \mathbb{R}^n$ except for the unit sphere $\{x = (x_1, \dots, x_n) | |x| = (x_1^2 + \dots + x_n^2)^{\frac{1}{2}} = 1\}$, and this extended hyperbolic space $(\{1\} \times \mathbb{R}^n, ds_K^2)$ will be denoted by K^n . Hence K^n consists of $\mathbb{H}^n, \mathbb{L}^n$, and the unit sphere.

Note that K^n is simply an affine coordinate chart of the *projective space* $\mathbb{R}P_H^n$ equipped with the singular metric composed of the usual hyperbolic metric coming from H_+^n (or H_-^n) and the negative of the Lorentz metric coming from S_1^n .

In this paper, it would be more convenient to consider the Euclidean unit sphere in $\mathbb{R}^{n+1}$ with the induced (singular) metric coming from this projective space as a double covering, and denote this model by $\mathbb{S}_H^n$. This *hyperbolic sphere model* $\mathbb{S}_H^n$ on the Euclidean sphere $\{x = (x_0, x_1, \dots, x_n) | x_0^2 + x_1^2 + \dots + x_n^2 = 1\}$ has three open parts: Two radial images of $H_\pm^n$, called the *hyperbolic part*, as two open disks on upper and lower hemisphere and the radial image of S_1^n , called the *Lorentzian part*, forms the remaining part which is diffeomorphic to $\mathbb{R}^1 \times \mathbb{S}^{n-1}$ (see Fig.2.1). Note that the image of S_1^1 is composed of two components.

In this section, we want to perceive this singular metric on $\mathbb{S}_H^n$ as an analytic continuation of the hyperbolic part. One way to do it is to view the metric as a limit of non-singular complex metric.

We define an ϵ -approximation of ds_K^2 as

$$ds_\epsilon^2 = \left(\frac{\sum x_i dx_i}{d_\epsilon^2 - |x|^2} \right)^2 + \frac{\sum dx_i^2}{d_\epsilon^2 - |x|^2},$$




Fig 2.1

where $d_\epsilon = 1 - \epsilon i$ with $\epsilon > 0$ and $i = \sqrt{-1}$. Then $ds_K^2 = \lim_{\epsilon \rightarrow 0} ds_\epsilon^2$.

Since the length and angle can be viewed as an 1-dimensional measure, we first study the volume (or measure) on $\mathbb{S}_H^n$ as a limit of the non-singular complex volume obtained from ds_ϵ^2 . Note that the volume form on K^n follows as

$$\begin{aligned} dV_K &= (\det(g_{ij}))^{\frac{1}{2}} dx_1 \wedge \cdots \wedge dx_n, \\ &= \frac{dx_1 \wedge \cdots \wedge dx_n}{(1 - |x|^2)^{\frac{n+1}{2}}}. \end{aligned}$$

The volume form for ds_ϵ^2 similarly will be given by

$$dV_\epsilon = \frac{d_\epsilon dx_1 \wedge \cdots \wedge dx_n}{(d_\epsilon^2 - |x|^2)^{\frac{n+1}{2}}}.$$

In the Kleinian model, for a set U in the unit disk H^n the volume of U will be simply given by

$$\text{vol}(U) = \int_U dV_K = \int_U \frac{dx_1 \wedge \cdots \wedge dx_n}{(1 - |x|^2)^{\frac{n+1}{2}}}$$

as far as the integral exists. Furthermore this integral is also obtained as $\lim_{\epsilon \rightarrow 0} \int_U dV_\epsilon$, since $|d_\epsilon - |x|^2| \geq |1 - |x|^2|$ and hence the Lebesgue dominated convergence theorem applies. We have the same conclusion for U lying solely in the Lorentzian part S_1^n . But when we consider the volume function for Lorentzian case, there is a sign problem. There is a natural consistent way of choosing a sign $\{\pm i, \pm 1\}$ for the volume computation in the Lorentzian part once $d_\epsilon = 1 - \epsilon i$ is chosen as our approximation, and the sign convention will be explained later.

Now if a subset U of K^n lies across the light cone, that is the boundary of $\mathbb{H}^n$, then the integral for $\text{vol}(U)$ does not make sense any more, and we want to define a volume of U as

$$\mu(U) = \lim_{\epsilon \rightarrow 0} \int_U \frac{d_\epsilon dx_1 \wedge \cdots \wedge dx_n}{(d_\epsilon^2 - |x|^2)^{\frac{n+1}{2}}} \quad (1)$$

whenever the limit exists. From now on we will call a Lebesgue measurable set U *μ -measurable* if $\mu(U)$ is defined as a finite value.

The actual computation of such integral doesn't seem to be easy. But we can show that $\mu(U)$ is equal to the integral of the singular volume form dV_K over U calculated in polar coordinates if it is interpreted appropriately for a nice class of the subsets U . One such case is when the integral can be considered as an analytic continuation in the radial direction in the polar coordinates. Consider the integral for $\text{vol}(U)$ as before.

$$\begin{aligned} \text{vol}(U) &= \int_U dV_K \\ &= \int_U \frac{dx_1 \cdots dx_n}{(1 - |x|^2)^{\frac{n+1}{2}}} \\ &= \int_{G^{-1}(U)} \frac{r^{n-1}}{(1 - r^2)^{\frac{n+1}{2}}} dr d\theta, \end{aligned}$$

where $G : (r, \theta) \mapsto (x_1, \dots, x_n)$ is the polar coordinates and $d\theta$ is the volume form of the Euclidean unit sphere $\mathbb{S}^{n-1}$. If $F(r) = \int_{G^{-1}(U) \cap \mathbb{S}^{n-1}(r)} d\theta$ is an analytic function of r , then we call such a subset U of K^n a *proper* set and the above integral becomes a 1-dimensional integral as follows.

$$\int_{G^{-1}(U)} \frac{r^{n-1}}{(1 - r^2)^{\frac{n+1}{2}}} dr d\theta = \int_a^b \frac{r^{n-1} F(r)}{(1 - r^2)^{\frac{n+1}{2}}} dr$$

In general this integral does not make sense and the most natural thing we can do instead is to define $\text{vol}(U)$ as the following contour integral

$$\text{vol}(U) := \int_\gamma \frac{r^{n-1} F(r)}{(1 - r^2)^{\frac{n+1}{2}}} dr \quad (2)$$

where γ is a contour from a to b for $a < 1 < b$ as depicted below. Here we will fix its contour type as clockwise around $z = 1$ once and for all throughout the paper.




Fig 2.2

Therefore we can compute the length of line segment on $\mathbb{S}_H^1$ by using the line integral (2). It is easy to see that

$$m(b) := \int_{0,\gamma}^b \frac{dr}{1-r^2} = \begin{cases} \frac{1}{2} \log \frac{1+b}{1-b}, & 0 \leq b < 1, \\ \frac{1}{2} \log \frac{b+1}{b-1} + \frac{\pi}{2}i, & 1 < b, \end{cases} \quad (3)$$

where the notation $\int_{a,\gamma}^b$ denote a line integral from a to b along the contour γ .

Now we can show that this contour integral in fact gives us a way of calculating $\mu(U)$.

Proposition 2.1 *For a proper set U in K^n , $\mu(U) = \text{vol}(U)$.*

Proof Using polar coordinates,

$$\begin{aligned} \mu(U) &= \lim_{\epsilon \rightarrow 0} \int_{G^{-1}(U)} \frac{d_\epsilon r^{n-1}}{(d_\epsilon^2 - r^2)^{\frac{n+1}{2}}} dr d\theta \\ &= \lim_{\epsilon \rightarrow 0} \int_a^b \frac{d_\epsilon r^{n-1} F(r)}{(d_\epsilon^2 - r^2)^{\frac{n+1}{2}}} dr \end{aligned}$$

for some $0 \leq a < 1 < b$. Then the equation $r^2 - d_\epsilon^2 = 0$ gives poles $r = \pm d_\epsilon = \pm(1 - \epsilon i)$, which, near 1, is located in the lower half of the complex plane $\mathbb{C}$. Hence the integral $\int_a^b$ is equal to $\int_\gamma$ for any contour γ from a to b lying in the upper half of $\mathbb{C}$. Now for such contour,

$$\mu(U) = \lim_{\epsilon \rightarrow 0} \int_\gamma \frac{d_\epsilon r^{n-1} F(r)}{(d_\epsilon^2 - r^2)^{\frac{n+1}{2}}} dr. \quad (4)$$

In particular, choose γ as depicted in Fig. 2.2 so that it goes around $z = 1$ through an upper semi-circle of radius δ in the clockwise direction. The upper semi-circle part of γ can be given by

$$\gamma(t) = 1 + \delta e^{(\pi-t)i}, \quad 0 \leq t \leq \pi.$$

Now for all sufficiently small $\epsilon < \epsilon_0$, choose δ such that $2\sqrt{\epsilon_0^4 + 4\epsilon_0^2} < \delta < 1$. Then we have

$$\begin{aligned} |d_\epsilon^2 - r^2| &= |-\epsilon^2 - 2\epsilon i + 1 - r^2| \geq |1 - r^2| - \sqrt{\epsilon^4 + 4\epsilon^2} \\ &\geq |1 - r| |1 + r| - \frac{\delta}{2} = \delta |2 + \delta e^{(\pi-t)i}| - \frac{\delta}{2} \geq \delta - \frac{\delta}{2} = \frac{\delta}{2}. \end{aligned}$$

This inequality gives us an uniform estimate of the integrand in (4) and completes the proof using Lebesgue dominated convergence theorem. $\square$

Remark 2.2 *Our choice of ϵ -approximation $d_\epsilon = 1 - \epsilon i$ is rather elaborate anticipating proposition 2.1. If we choose $\tilde{d}_\epsilon = 1 + \epsilon i$ instead, then $\tilde{\mu}(U) = \lim_{\epsilon \rightarrow 0} \int_U d\tilde{V}_\epsilon$ will be the same as the $\tilde{\text{vol}}(U)$ which is given as in (1) where the contour type is given as counterclockwise around $z = 1$, i.e., lower going at $z = 1$. The total length of $\mathbb{S}_H^1$ is $2\pi i$ for the choice of d_ϵ , and $-2\pi i$ for the choice of $\tilde{d}_\epsilon$.*

To determine the various geometric quantities like the integration for arc length on $\mathbb{S}_H^n$, the notion of norms of vectors is essential. From the sign change of the metric on the pseudo-sphere S_1^n , the norm of tangent vector $x_p = (x_0, x_1, \dots, x_n)$ at a Lorentzian part point is given by

$$\|x_p\|^2 = -(-x_0^2 + x_1^2 + \dots + x_n^2),$$

and we should determine the sign of $\|x_p\|$ among ± 1 and $\pm i$.

On the 1-dimensional hyperbolic sphere $\mathbb{S}_H^1$, we can determine the sign by looking at the sign of arc-length which can be calculated by the formula (3). Since the function $m(b)$ in the formula (3) is decreasing on $b > 1$, the sign of arc-length becomes negative value outside of $\mathbb{H}^1$ (see Fig. 2.3). This gives us (-1) as the right choice of the sign of $\|x_p\|$ for the vectors in the radial direction at a Lorentzian point.

For the sign for the vectors normal to the radial direction, we need the following argument. It is not hard to check that the clockwise contour integral of the volume form gives sign $-i^{n-1}$ for Lorentzian part. Indeed for the volume integral in the Lorentzian part will be given as

$$\int_a^b f(r)dr = \int_{0,\gamma}^b f(r)dr - \int_{0,\gamma}^a f(r)dr, \quad 1 < a < b,$$

integrated along the clockwise contour γ . Here the integrand $f(r) = \frac{r^{n-1}F(r)}{(1-r^2)^{\frac{n+1}{2}}}$ can be written as $f(r) = \frac{g(r)}{(1-r^2)^{\frac{n+1}{2}}}$ with $g(r) \geq 0$ for $r \geq 0$. The real axis for $r > 1$ will have the same sign under the map $f(r)$ and the sign can be traced using the contour γ which can be parametrized as $r = 1 + \delta e^{(\pi-t)i}, 0 \leq t \leq \pi$, near $r = 1$. Then $\frac{1}{(1-r^2)^{\frac{n+1}{2}}} = a e^{\frac{n+1}{2}ti} (2 + \delta e^{(\pi-t)i})^{-\frac{n+1}{2}}, a = \delta^{-\frac{n+1}{2}} > 0$, the term $(2 + \delta e^{(\pi-t)i})^{-\frac{n+1}{2}}$ doesn't give an effect on the sign at $t = \pi$ and thus the sign at $t = \pi$ will be $e^{\frac{n+1}{2}\pi i} = i^{n+1} = -i^{n-1}$. Hence on the 2-dimensional hyperbolic sphere $\mathbb{S}_H^2$, the volume for Lorentzian part has the sign $-i$ and thus the consistent choice of sign for the normal direction is i (see Fig. 2.3). Now since the dimension of the normal direction is $n - 1$ with sign i and radial direction with sign -1 , the volume has sign $(-1) \cdot i^{n-1} = -i^{n-1}$, which confirms again conceptually our volume sign for the Lorentzian part.

It is easy to see that the light cone at a point in the Lorentzian part is precisely the cone which is tangent to $\partial\mathbb{H}^n$ (see Fig. 2.3). Recall that our metric is the negative of the metric of the pseudo-sphere S_1^n and hence it has signature $(+, -, \dots, -)$, which implies that $\|x_p\|^2 > 0$ inside the light cone (the shaded region in Fig. 2.3) and $\|x_p\|^2 < 0$ for normal directions. We can see this in the Minkowski space by looking at the tangent vector x_p at $p \in S_1^n$ which lies inside and outside of the light cone at p respectively.

We summarize the above discussion as the following convention.

Convention *A tangent vector on the hyperbolic part on $\mathbb{S}_H^n$ has a positive real norm, and a tangent vector on the Lorentzian part on $\mathbb{S}_H^n$ has a negative real, zero, or positive pure imaginary norm depending on whether it is timelike, null, or spacelike respectively.*

In [16], we can see a complex distance extending the cross ratio to the exterior of the hyperbolic space. And we can easily check that the definition using cross ratio and our




Fig 2.3

definition for distance and angle coincide. Our definition of angle will be explained in Section 3.

We can see one of the similarities between the Euclidean sphere $\mathbb{S}^n$ and the hyperbolic sphere $\mathbb{S}_H^n$ in the following theorem.

Theorem 2.3 $vol(\mathbb{S}_H^n) = i^n \cdot vol(\mathbb{S}^n)$.

Proof Putting $U = \mathbb{S}_H^n$ in the formula for $vol(U)$ above, we have

$$vol(\mathbb{S}_H^n) = 2 \left(\int_{0,\gamma}^{\infty} \frac{r^{n-1}}{(1-r^2)^{\frac{n+1}{2}}} dr \right) vol(\mathbb{S}^{n-1}). \quad (5)$$

Similarly, we get the volume of the standard sphere by the radial projection from $\mathbb{S}^n$ to $\{1\} \times \mathbb{R}^n$, and that is represented as

$$vol(\mathbb{S}^n) = 2 \left(\int_0^{\infty} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt \right) vol(\mathbb{S}^{n-1}). \quad (6)$$

To compare (5) and (6), we need to define some contours (see Fig. 2.4),

$$C_N : z_1(t) = \begin{cases} t, & 0 \leq t \leq 1 - \epsilon, \\ 1 + \epsilon e^{\frac{i(1-t)\pi}{\epsilon}}, & 1 - \epsilon \leq t \leq 1, \\ t + \epsilon, & 1 \leq t \leq N - \epsilon, \end{cases}$$

$$C'_N : z_2(t) = -iz_1(t),$$

$$C''_N : z_3(t) = Ne^{-it}, \quad 0 \leq t \leq \frac{\pi}{2}.$$




Fig 2.4

For a given clockwise contour, we get

$$\begin{aligned}
& \int_{0,\gamma}^{\infty} \frac{r^{n-1}}{(1-r^2)^{\frac{n+1}{2}}} dr \\
&= \lim_{N \rightarrow \infty} \int_{C_N} \frac{r^{n-1}}{(1-r^2)^{\frac{n+1}{2}}} dr \\
&= \lim_{N \rightarrow \infty} i^n \int_{C'_N} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt \quad r = it \\
&= \lim_{N \rightarrow \infty} i^n \left(\int_{C_N} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt + \int_{C''_N} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt \right) \\
&= \lim_{N \rightarrow \infty} i^n \int_{C_N} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt \\
&= i^n \int_0^{\infty} \frac{t^{n-1}}{(1+t^2)^{\frac{n+1}{2}}} dt.
\end{aligned}$$

From (5),(6), and the above relation, we deduce that $\text{vol}(\mathbb{S}_H^n) = i^n \text{vol}(\mathbb{S}^n)$. $\square$

Remark 2.4 If we change the contour type of the integral (2), we have different relation between $\text{vol}(\mathbb{S}_H^n)$ and $\text{vol}(\mathbb{S}^n)$. If the contour is counterclockwise, in this case we should replace d_ϵ to $\tilde{d}_\epsilon = 1 + \epsilon i$, then we have $\text{vol}(\mathbb{S}_H^n) = (-i)^n \text{vol}(\mathbb{S}^n)$.

Remark 2.5 For various kinds of contour types, we easily deduce the following two formulas,

$$\begin{aligned}
\text{vol}(\mathbb{S}_H^{2k-1}) &\equiv i^{2k-1} \text{vol}(\mathbb{S}^{2k-1}) \quad (\text{mod } 2i^{2k-1} \text{vol}(\mathbb{S}^{2k-1})), \\
\text{vol}(\mathbb{S}_H^{2k}) &= i^{2k} \text{vol}(\mathbb{S}^{2k}).
\end{aligned}$$

Above formulas say that the total volume of even dimensional model has unique value for any contour but odd dimensional model has infinitely many values for various types of contours.

For basic two contour types, we can go through into well established theory by using of d_ϵ and $\tilde{d}_\epsilon$.

In the hyperbolic sphere $\mathbb{S}_H^n$, the hyperbolic part has $+\infty$ volume and the Lorentzian part has $+\infty i^{n+1}$ ($+\infty(-i)^{n+1}$ for $\tilde{d}_\epsilon$ case) volume. So the fact that the volume of $\mathbb{S}_H^n$ is $i^n \text{vol}(\mathbb{S}^n)$ gives the following nonsense equality.

$$\infty + \infty \cdot i^{n+1} = \text{vol}(\mathbb{S}^n) \cdot i^n$$

or

$$\infty + \infty \cdot (-i)^{n+1} = \text{vol}(\mathbb{S}^n) \cdot (-i)^n, \text{ for } \tilde{d}_\epsilon \text{ case.}$$

But the nonsense equality does not come true in a suitable measure theory, hence we need not worry about that. In Section 3, we can see the suitable and nice finitely additive complex measure theory for the spaces K^n , $\mathbb{R}P_H^n$, and $\mathbb{S}_H^n$.

3 Invariant property and measure theory on the extended model

Let us study the invariance of μ -measurable set in $\mathbb{R}P_H^n$ or $\mathbb{S}_H^n$. Here the isometry group of $\mathbb{S}_H^n$ is $O(n, 1)$ and we identify $PO(n, 1)$ as the subgroup of $O(n, 1)$ leaving H_+^n invariant. If U is a measurable set in the hyperbolic part or in the Lorentzian part, then the μ -measure is simply the usual hyperbolic or Lorentzian volume (with appropriate sign) and hence the invariance is obvious. But if we consider U lying across the $\partial\mathbb{H}^n$, the boundary of the hyperbolic part, then the invariance seems to be a subtle problem. Even if we do not know the invariance for general μ -measurable set holds, we are able to show the invariance for a region with piecewise analytic boundary transversal to $\partial\mathbb{H}^n$. More precisely, we define as follows.

Definition 3.1 *An n -dimensional region U of the extended hyperbolic space $\mathbb{S}_H^n$ is called a region with piecewise analytic boundary transversal to $\partial\mathbb{H}^n$ if it is bounded by finitely many analytic sets, $A_i := \{x | f_i(x) = 0, f_i \text{ is an analytic function}\}$, $i = 1, 2, \dots, k$, which are transversal to $\partial\mathbb{H}^n$ such that for each $p \in U \cap \partial\mathbb{H}^n$, p belongs to one of the bounding analytic sets but at most two such sets A_i and $A_{i+1} \pmod k$ with the property that $\nabla f_i(p)$, $\nabla f_{i+1}(p)$ and the normal vector to $\partial\mathbb{H}^n$ are linearly independent.*

Proposition 3.2 *Let U be a region with piecewise analytic boundary transversal to $\partial\mathbb{H}^n$ in the extended hyperbolic space. Then $\mu(U)$ has a well-defined finite value, i.e., U is μ -measurable, and $\mu(g(U)) = \mu(U)$ for each $g \in PO(n, 1)$.*

Proof We want to show first that $F_U(r) = \int_{G^{-1}(U) \cap S^{n-1}(r)} d\theta$ and $F_{gU}(r)$ are real analytic function of r near $r = 1$, i.e., U and gU are proper sets, (see the previous section for the notations and definition) so that U and gU have well-defined finite values.

We decompose a neighborhood of U near $r = 1$ appropriately so that each piece has a box coordinate under a real analytic coordinate change as in Figure 3.1. If we denote




Fig 3.1

two analytic boundaries in V_m by $f_m(x_1, \dots, x_{n-1}, r) = 0$ and $f_{m+1}(x_1, \dots, x_{n-1}, r) = 0$, then we can rewrite these as $x_1 = g_m(x_2, \dots, x_{n-1}, r)$ and $x_1 = g_{m+1}(x_2, \dots, x_{n-1}, r)$ by the (analytic) implicit function theorem. From the condition that $\nabla f_m(x_1, \dots, x_{n-1}, 1)$ and $\nabla f_{m+1}(x_1, \dots, x_{n-1}, 1)$ are not parallel, we may assume $\frac{\partial}{\partial x_2} g_m \neq \frac{\partial}{\partial x_2} g_{m+1}$, and then again by the (analytic) implicit function theorem, we can write the intersection of two analytic sets, $g_m(x_2, \dots, x_{n-1}, r) = g_{m+1}(x_2, \dots, x_{n-1}, r)$, as $x_2 = h(x_3, \dots, x_{n-1}, r)$.

Now the function $F_U(r)$ for the shaded region U in V_m in Fig. 3.1 will be given by

$$F_U(r) = \int_{a_{n-1}}^{b_{n-1}} \cdots \int_{a_3}^{b_3} \int_{a_2}^{h(x_3, \dots, x_{n-1}, r)} \int_{a_1}^{g_m(x_2, \dots, x_{n-1}, r)} f(x) dx_1 dx_2 \cdots dx_{n-1} \\ + \int_{a_{n-1}}^{b_{n-1}} \cdots \int_{a_3}^{b_3} \int_{h(x_3, \dots, x_{n-1}, r)}^{g_{m+1}(x_2, \dots, x_{n-1}, r)} \int_{a_1}^{g_{m+1}(x_2, \dots, x_{n-1}, r)} f(x) dx_1 dx_2 \cdots dx_{n-1}.$$

Since the function h, g_m, g_{m+1} , and f are analytic, $F_U(r)$ is analytic as desired. Similarly $F_{gU}(r)$ is also analytic.

Next we show that $\mu(U) = \mu(gU)$. Let γ be a contour from a to b in Fig. 2.2 and let $B_r = G^{-1}(U) \cap S^{n-1}(r)$. Then

$$\mu(U) = \lim_{\epsilon \rightarrow 0} \int_U dV_\epsilon = \lim_{\epsilon \rightarrow 0} \int_a^b \frac{r^{n-1} F_U(r)}{(d_\epsilon^2 - r^2)^{\frac{n+1}{2}}} dr \\ = \lim_{\epsilon \rightarrow 0} \int_\gamma \frac{r^{n-1} F_U(r)}{(d_\epsilon^2 - r^2)^{\frac{n+1}{2}}} dr = \int_\gamma \frac{r^{n-1} F_U(r)}{(1 - r^2)^{\frac{n+1}{2}}} dr = \int_\gamma \int_{B_r} dV_0.$$

Here the third equality holds since the pole of the integrand near 1 has a negative imaginary part. The fourth equality follows from $|d_\epsilon^2 - r^2| \geq \frac{\delta}{2}$ as in the proof of Proposition 2.1 so that the integrand is uniformly bounded for all sufficiently small $\epsilon > 0$ and we can apply Lebesgue dominated convergence theorem.

Now if we can show for $\mu(gU)$ similarly as follows, then the proof will be completed.

$$\begin{aligned}\mu(gU) &= \lim_{\epsilon \rightarrow 0} \int_{gU} dV_\epsilon = \lim_{\epsilon \rightarrow 0} \int_U g^*(dV_\epsilon) = \lim_{\epsilon \rightarrow 0} \int_a^b \int_{B_r} g^*(dV_\epsilon) \\ &= \lim_{\epsilon \rightarrow 0} \int_\gamma \int_{B_r} g^*(dV_\epsilon) = \int_\gamma \int_{B_r} g^*(dV_0) = \int_\gamma \int_{B_r} dV_0.\end{aligned}$$

Here g is an isometry and hence $g^*(dV_0) = dV_0$, but notice that g does not preserve ϵ -volume form dV_ϵ and we need several steps as before. We have to check 4th and 5th equality and all others are obvious.

For $g \in PO(n, 1)$, if we let $dV_\epsilon = f_\epsilon(r, \theta) d\theta dr$, then

$$g^*(dV_\epsilon) = (f_\epsilon \circ g)(\det g') d\theta dr =: h_\epsilon(r, \theta) d\theta dr.$$

Since g is an isometry, g obviously preserves the orientation of real r -axis. When we complexify everything, as an analytic map g preserves the orientation of complex r -axis and leaves real r -axis invariant being a real map. Hence the pole of h_ϵ also lies below the real axis as f_ϵ does.

Now for the 5th equality, note that dV_ϵ and $g^*(dV_\epsilon)$ differ only by Jacobian determinant for their values. Since we already know that dV_ϵ is uniformly bounded on its domain of integral $\gamma \times B_r$ so is $g^*(dV_\epsilon)$ and we can apply Lebesgue dominated convergence theorem to obtain the equality. $\square$

Remark 3.3 *In fact, we can generalize this proposition to a region with more general type boundaries than analytic boundaries. (See [5].)*

Let $\mathcal{F}$ be the collection of μ -measurable subsets of K^n . Then it is obvious that $\mu : \mathcal{F} \rightarrow \mathbb{C}$ is a finitely additive set function, i.e., for $E_1, E_2 \in \mathcal{F}$ and disjoint union $E_1 \amalg E_2$, we have $\mu(E_1 \amalg E_2) = \mu(E_1) + \mu(E_2)$ so $E_1 \amalg E_2 \in \mathcal{F}$ (this result trivially generalizes to that if two sets among $E_1, E_2, E_1 \amalg E_2$ are μ -measurable, then the other is μ -measurable).

It would be nice if $\mu : \mathcal{F} \rightarrow \mathbb{C}$ is a finitely additive measure on an algebra $\mathcal{F}$. But unfortunately $\mathcal{F}$ is not an algebra, as we see in the following example.

Consider a cone E_1 over a small disk B on $\partial\mathbb{H}^n$ with the vertex at the origin of $\mathbb{H}^n$, where the origin is the point $\{1\} \times \{0\} \in \{1\} \times \mathbb{R}^n = K^n$. Let g be a rotation such that $g(E_1 \cap \mathbb{H}^n)$ is disjoint from $E_1 \cap \mathbb{H}^n$, and let $E_2 = g(E_1 \cap \mathbb{H}^n) \cup (E_1 \cap \mathbb{L}^n)$. Then E_1 and E_2 are clearly μ -measurable sets and by Theorem 2.3, for $i = 1, 2$,

$$\begin{aligned}\mu(E_i) &= \text{vol}(E_i) = \left(\int_{0, \gamma}^{\infty} \frac{r^{n-1}}{(1-r^2)^{\frac{n+1}{2}}} dr \right) \text{vol}(B) \\ &= \frac{i^n \text{vol}(\mathbb{S}^n)}{2 \text{vol}(\mathbb{S}^{n-1})} \text{vol}(B) = \frac{i^n \sqrt{\pi}}{2} \frac{\Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \text{vol}(B).\end{aligned}$$

But $E_1 \cap E_2$ and $E_1 \cup E_2$ have infinite volume, hence not μ -measurable. This example looks rather artificial, but we will show that the problem of finding a useful large enough algebra in $\mathcal{F}$ is a very sensitive problem in its nature.

The simplest type of μ -measurable sets in $\mathbb{H}^n$ (or in $\mathbb{L}^n$) is bounded regions not touching the $\partial\mathbb{H}^n$, because the μ -measurable sets in $\mathbb{H}^n$ (or in $\mathbb{L}^n$) is exactly the same as the original positive volume measure in $\mathbb{H}^n$ (or in $\mathbb{L}^n$ up to sign). Also the cone region with vertex at the origin of $\mathbb{H}^n$ is μ -measurable. Our first attempt to finding an algebra in $\mathcal{F}$ is the one generated by sets of the above three types.

Let $\mathcal{U}_h$ (resp. $\mathcal{U}_l$) be the collection of μ -measurable sets in $\mathbb{H}^n$ (resp. $\mathbb{L}^n$) and $\mathcal{U}_c$ be the collection of μ -measurable cones with vertex at the origin of $\mathbb{H}^n$. We assume that $\emptyset \in \mathcal{U}_c$. Let $\mathcal{M}$ be the smallest algebra containing $\mathcal{U}_h, \mathcal{U}_l$ and $\mathcal{U}_c$, so that $\mathcal{M}$ contains $\mathcal{U}_d := \{V \mid V = U_2 - U_1, U_2 \in \mathcal{U}_c \text{ and } U_1 = U_h \cup U_l, \text{ where } U_h \in \mathcal{U}_h \text{ and } U_l \in \mathcal{U}_l\}$.

Proposition 3.4 (1) Let $\mathcal{M}' = \{V_1 \cup V_2 \mid V_2 \in \mathcal{U}_d \text{ and } V_1 = V_h \cup V_l, \text{ where } V_h \in \mathcal{U}_h \text{ and } V_l \in \mathcal{U}_l\}$ and $\mathcal{M}''$ be the subcollection of $\mathcal{M}'$ with the condition $V_1 \cap V_2 = \emptyset$. Then we have $\mathcal{M} = \mathcal{M}' = \mathcal{M}''$.

(2) For any $U \in \mathcal{M}$, U can be written as a disjoint union $U = U_h \amalg U_l \amalg U_d$, where $U_h \in \mathcal{U}_h$, $U_l \in \mathcal{U}_l$ and $U_d \in \mathcal{U}_d$. Hence the collection $\mathcal{M}$ is a μ -measurable algebra.

Proof An element of $\mathcal{M}'$ can be written as $V_1 \cup (U_2 - U_1)$ and the proof that $\mathcal{M}' = \mathcal{M}''$ follows readily from the general set identity:

$$V_1 \cup (U_2 - U_1) = (V_1 - U_2) \amalg (U_2 - (U_1 - V_1)).$$

We want to show that $\mathcal{M} = \mathcal{M}'$. Since $\mathcal{M}' \subset \mathcal{M}$ and $\mathcal{U}_h, \mathcal{U}_l, \mathcal{U}_c \subset \mathcal{M}'$, it suffices to show that $\mathcal{M}'$ is an algebra. The whole space $K^n \in \mathcal{M}'$ since it is a cone. It is easy to show that $V \in \mathcal{M}'$ implies $V^c \in \mathcal{M}'$. Indeed this follows from the general set identity:

$$(V_1 \cup (U_2 - U_1))^c = (U_1 - V_1) \cup (U_2^c - V_1).$$

Note that the complement of a cone is also a cone.

Finally, let's show that $\mathcal{M}'$ is closed under the union operation. Write

$$(V_1 \cup (U_2 - U_1)) \cup (V'_1 \cup (U'_2 - U'_1)) = (V_1 \cup V'_1) \cup ((U_2 - U_1) \cup (U'_2 - U'_1)).$$

and observe the set identity:

$$(U_2 - U_1) \cup (U'_2 - U'_1) = (U_2 \cup U'_2) - ((U_1 - U'_2) \cup (U'_1 - U_2) \cup (U_1 \cap U'_1)).$$

Now (2) follows from (1) trivially. $\square$

In dimension 2, $\mathcal{M}$ gives us a fairly large class of μ -measurable sets. We know that a cusp in $\mathbb{H}^2$ has a finite area. The same thing holds for a cusp in $\mathbb{L}^2$. This can be shown easily by a direct computation of the integral in the plane K^2 , or by looking at a configuration of three cones depicted in Fig. 3.2, if we notice that any cone with vertex in $\mathbb{H}^2$ has a finite area by proposition 3.2.

These observations and that μ -measure is essentially a positive measure in $\mathbb{H}^n$ and $\mathbb{L}^n$ lead us immediately to conclude that a domain transversal to $\partial\mathbb{H}^2$ belongs to $\mathcal{M}$ without using Proposition 3.2 (see Fig. 3.3).




Fig 3.2




Fig 3.3

But if the dimension is greater than or equal to 3, the algebra $\mathcal{M}$ becomes a small collection which doesn't even contain the class of cones with vertex in $\mathbb{H}^n$, whose volume, as we already know, are finite.

Indeed let's consider the following two cones in $\mathbb{R}^n = K^3$.

$$C_1 : z \geq \frac{3}{4}r$$

$$C_2 : z \geq \frac{1}{3}(r+1)$$

Then $C_1 \in \mathcal{M}$ by the definition of $\mathcal{M}$, but we claim $C_2 \notin \mathcal{M}$ by showing that the volume of $E = C_2 - C_1$ is infinite by direct computation.

$$\begin{aligned} \mu(E) &= \iiint_E \frac{dxdydz}{(1-x^2-y^2-z^2)^2} \\ &= \int_{\frac{3}{5}}^{\delta} \int_0^{2\pi} \int_{\frac{4}{3}z}^{3z-1} \frac{rdrd\theta dz}{(1-z^2-r^2)^2} \\ &= \pi \int_{\frac{3}{5}}^{\delta} \frac{1}{z - \frac{3}{5}} \left(\frac{9}{25} \frac{1}{z + \frac{3}{5}} - \frac{1}{10z} \right) dz. \end{aligned}$$

This integral diverges since the function in parenthesis in the integrand has a positive lower bound on $[\frac{3}{5}, \delta]$.




Fig 3.4

Even though the collection $\mathcal{M}$ does not contain natural domains we want in dimensions greater than or equal to 3, we can construct sufficiently large and very important collections $\mathcal{H}$ and $\mathcal{H}'$ in such dimensions.

Proposition 3.5 *Let $\mathcal{H}$ be the algebra in K^n (resp. $\mathbb{S}_H^n$) generated by half spaces (resp. hemisphere) not tangent to $\partial\mathbb{H}^n$. Then the collection $\mathcal{H}$ is a μ -measurable algebra.*

Proof To show μ -measurability of $U \in \mathcal{H}$, it suffices to consider only μ -measurability of the region $U \cap \{1 - \delta \leq r \leq 1 + \delta\}$ for a small $\delta > 0$ since the remaining parts are completely contained in either $\mathbb{H}^n$ or $\mathbb{L}^n$.




Fig 3.5

When the dimension $n = 2$, by virtue of Proposition 3.2, the only exceptional case we have to worry are the cone types regions K, L, M, N given as in Fig 3.5-(ii). We know that the hyperbolic cusp M and the Lorentzian cusp L have finite areas.

The key observation about the μ -measurability we are using over and over in the following argument is that if two of A, B and $A \cup B$ are μ -measurable, so is the remaining one. For

instance, K is μ -measurable since L and $K \cup L$ are μ -measurable. In this way all these four types of region are μ -measurable.

When $n = 3$, we have to consider several exceptional case.

Case 1. When two planes transversal to $\partial\mathbb{H}^3$ meet at a line which is tangent to $\partial\mathbb{H}^3$: (See Fig. 3.5-(iii).) Again in this case, we have four regions K, L, M, N as before and we can show that the region L contained in the Lorentzian part has a finite volume by direct computations (see APPENDIX). Then the other regions have finite volumes by the same argument as the case $n = 2$ using the key observation.

Case 2. When three transversal planes meet at a point on $\partial\mathbb{H}^3$: In this case, we obtain 8 regions from these 3 planes. At least one of these regions is contained in the Lorentzian part and this is contained in an L -type region considered in Case 1. This implies that it has a finite volume since μ -measure on the Lorentzian part is essentially a positive measure. Then a neighboring octant has a finite volume by the key observation and Proposition 3.2. Hence again their neighboring octants have finite volumes by the same reasoning and so forth.

Case 3. When four or more transversal planes meet at a point on $\partial\mathbb{H}^3$: Fig. 3.5-(iv) shows a region (shaded one) bounded by 4 transversal planes. This region is the difference of two regions of the type considered in Case 2 and hence is of finite volume again by the key observation. Now finiteness of volume of a region bounded by many planes follows by induction.

If $n \geq 4$, we have more exceptional cases but still we have the same conclusion by the same argument and observations since we can start with a piece in the Lorentzian part which has a finite volume by APPENDIX. $\square$

Proposition 3.6 *Let $\mathcal{H}'$ be the smallest algebra in K^n or $\mathbb{S}_H^n$ containing $\mathcal{U}_h, \mathcal{U}_l$, and $\mathcal{H}$. Then the collection $\mathcal{H}'$ is μ -measurable algebra.*

Proof Use the same method shown in Proposition 3.4. $\square$

In dimension 2, the collection $\mathcal{M}$ is strictly larger than $\mathcal{H}'$, i.e., $\mathcal{H}' \subsetneq \mathcal{M}$. We will leave the proof that $\mathcal{H}' \subsetneq \mathcal{M}$ as an easy exercise for readers.

4 Lengths and angles on the extended hyperbolic space

We denote the distance between two points A and B in the extended hyperbolic space $\mathbb{S}_H^n$ as $d_H(A, B)$. Let's first discuss the distance between two points on $\mathbb{S}_H^1$. In this case, the ϵ -metric and the volume form are given as $ds_\epsilon^2 = \frac{d_\epsilon^2 dx_1^2}{(d_\epsilon^2 - x_1^2)^2}$ and $dV_\epsilon = \frac{d_\epsilon dx_1}{d_\epsilon^2 - x_1^2} = ds_\epsilon$ on its affine chart K^1 . If we let the affine coordinates of two points A and B of $\mathbb{S}_H^1$ be $x_1 = a$ and $x_1 = b$ respectively, then the μ -measure of the line segment $l = [a, b]$ is

$$\mu(l) = \lim_{\epsilon \rightarrow 0} \int_l ds_\epsilon = \lim_{\epsilon \rightarrow 0} \int_a^b \frac{d_\epsilon dx_1}{d_\epsilon^2 - x_1^2},$$

and this is to be the distance $d_H(A, B)$ of A and B . For instance, if A and B are symmetric with respect to the light cone $x_0 = x_1$ in $\mathbb{R}^{1,1}$ as in Fig. 4.1. (i.e., A and B as vectors of

$\mathbb{R}^{1,1}$ are perpendicular), then their affine coordinates are $a(< 1)$ and $\frac{1}{a}$, and the distance will be $\frac{\pi}{2}i$ by formula (3). The distance between isometric images A' and B' of A and B will be again $\frac{\pi}{2}i$ being symmetric, and hence $d_H(B, B') = -d_H(A, A')$ in Fig. 4.1.




Fig 4.1

To discuss the distance between two points in $\mathbb{S}_H^n$ in general, it suffices to consider on $\mathbb{S}_H^2$. It is natural to define a distance between two points on $\mathbb{S}_H^2$ by the μ -measure of the geodesic line segment connecting these two points.

For actual computations, it would be convenient to divide into the following 3 cases. For the case when the geodesic connecting two points meet $\partial\mathbb{H}^2$ transversely, we may assume that these two points lie on $\mathbb{S}_H^1 = \mathbb{S}_H^2 \cap \{x|x_2 = 0\}$ by an isometry and can handle as discussed above.

For the case when the geodesic line connecting these two points does not intersect $\partial\mathbb{H}^2$, we can send this line to the equator ($= \mathbb{S}_H^2 \cap \{x|x_0 = 0\}$) of $\mathbb{S}_H^2$ by an isometry, and hence the distance becomes i times the distance on the standard Euclidean unit circle.

The remaining case is when the line is tangent to $\partial\mathbb{H}^2$. On K^2 , if we restrict the Kleinian ϵ -metric to the line given by $x_2 = k$, then we have $ds_\epsilon^2 = \frac{(d_\epsilon^2 - k^2)dx_1^2}{(d_\epsilon^2 - k^2 - x_1^2)^2}$ and the length of the line segment connecting $(0, k)$ and (a, k) for $a > 0$ will be given by $\lim_{\epsilon \rightarrow 0} \int_0^a \frac{(d_\epsilon^2 - k^2)^{\frac{1}{2}} dx_1}{d_\epsilon^2 - k^2 - x_1^2}$. Since the length of the line segment in $\mathbb{H}^2$ is positive, we have to choose one whose real part is positive among the two possible values of $(d_\epsilon^2 - k^2)^{\frac{1}{2}}$ when $0 < k < 1$. By continuity we choose $c_\epsilon = (d_\epsilon^2 - 1)^{\frac{1}{2}}$ with $Re((d_\epsilon^2 - 1)^{\frac{1}{2}}) > 0$ (or $Im((d_\epsilon^2 - 1)^{\frac{1}{2}}) < 0$). Now the length of the line segment connecting $(0, 1)$ and $(a, 1)$ for $a > 0$ will be

$$\begin{aligned} \lim_{\epsilon \rightarrow 0} \int_0^a \frac{(d_\epsilon^2 - 1)^{\frac{1}{2}} dx_1}{d_\epsilon^2 - 1 - x_1^2} &= \lim_{\epsilon \rightarrow 0} \int_0^a \frac{c_\epsilon dx_1}{c_\epsilon^2 - x_1^2} = \lim_{\epsilon \rightarrow 0} \frac{1}{2} \log \frac{c_\epsilon + x_1}{c_\epsilon - x_1} \Big|_0^a \\ &= \lim_{\epsilon \rightarrow 0} \frac{1}{2} \log \frac{c_\epsilon + a}{c_\epsilon - a} = \lim_{\epsilon \rightarrow 0} \frac{1}{2} \log \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}} = \lim_{\epsilon \rightarrow 0} \left(\frac{1}{2} \log \frac{r_1}{r_2} + \frac{i(\theta_1 - \theta_2)}{2} \right) = \frac{\pi}{2}i \end{aligned}$$

Now from the finite additivity of μ -measure, we conclude as follows.

Lemma 4.1 *For a point x lying on $\partial\mathbb{H}^2$ and a dual geodesic $x^\perp$, the lengths of the line segments in $x^\perp$ are defined by*

$$\begin{aligned} d_H(w, y) &= 0, & \text{if } w, y \text{ are in the same side with respect to } x, \\ d_H(y, z) &= \pi i, & \text{if } y, z \text{ are in the opposite sides with respect to } x, \\ d_H(x, y) &= d_H(x, z) = \frac{\pi}{2}i. \end{aligned}$$

See the following Fig. 4.2.




Fig 4.2

In all of these discussions, we in fact have to show that 1-dimensional μ -measure is invariant under isometry since the hyperbolic isometry does not preserve ϵ -metric. We will show this in the following Theorem 4.3 below.

We conclude the following theorem from the above discussion.

Theorem 4.2 *The total length, i.e., 1-dimensional μ -measure, of any great circle in $\mathbb{S}_H^n$ is $2\pi i$.*

The intersection of $(k + 1)$ -dimensional subspace of $\mathbb{R}^{n,1}$ with $\mathbb{S}_H^n$ is a totally geodesic k -dimensional subspace of $\mathbb{S}_H^n$ and the ϵ -metric on this k -dimensional subspace induced from that of $\mathbb{S}_H^n$ gives rise to a k -dimensional ϵ -volume form and a k -dimensional μ -measure as a limit.

Theorem 4.3 *A k -dimensional region with piecewise analytic boundary contained in a k -dimensional totally geodesic subspace transversal to $\partial\mathbb{H}^n$ has a finite k -dimensional measure which is invariant under the isometry action.*

Proof Let U be a k -dimensional region contained in a geodesic sphere S^k . Here we assume S^k is transversal to $\partial\mathbb{H}^n$ and we will discuss the case when S^k is tangent to $\partial\mathbb{H}^n$ in the following remark.

If U does not intersect $\partial\mathbb{H}^n$, then the theorem is clear and we assume U intersect $\partial\mathbb{H}^n$. It suffices to show that $\mu_k(U)$ is finite and $\mu_k(U) = \mu_k(gU)$ for any $g \in PO(n, 1)$ when U is

contained in $l^k := \mathbb{S}_H^k = \mathbb{S}_H^n \cap \{x | x_{k+1} = \dots = x_n = 0\}$. In this case $\mu_k(U)$ is given as follows as in Proposition 3.2 and hence finite.

$$\begin{aligned}\mu_k(U) &= \lim_{\epsilon \rightarrow 0} \int_U dV_\epsilon^{l^k} = \lim_{\epsilon \rightarrow 0} \int_a^b \frac{r^{k-1} F(r)}{(d_\epsilon^2 - r^2)^{\frac{k+1}{2}}} dr \\ &= \lim_{\epsilon \rightarrow 0} \int_\gamma \frac{r^{k-1} F(r)}{(d_\epsilon^2 - r^2)^{\frac{k+1}{2}}} dr = \int_\gamma \frac{r^{k-1} F(r)}{(1 - r^2)^{\frac{k+1}{2}}} dr = \int_\gamma \int_{B_r} dV_0^{l^k},\end{aligned}$$

when $dV_\epsilon^{l^k}$ is the k -dimensional ϵ -volume form on $l^k := \mathbb{S}_H^k$.




Fig 4.3

Now we show that $\mu_k(gU) = \int_\gamma \int_{B_r} dV_0^{l^k}$ for $g \in PO(n, 1)$ as before by checking the following steps.

$$\begin{aligned}\mu_k(gU) &= \lim_{\epsilon \rightarrow 0} \int_{gU} dV_\epsilon^{m^k} = \lim_{\epsilon \rightarrow 0} \int_U g^*(dV_\epsilon^{m^k}) = \lim_{\epsilon \rightarrow 0} \int_a^b \int_{B_r} g^*(dV_\epsilon^{m^k}) \\ &= \lim_{\epsilon \rightarrow 0} \int_\gamma \int_{B_r} g^*(dV_\epsilon^{m^k}) = \int_\gamma \int_{B_r} g^*(dV_0^{m^k}) = \int_\gamma \int_{B_r} dV_0^{l^k}.\end{aligned}$$

But the proofs of these are exactly same if we notice that the k -dimensional ϵ -volume form $dV_\epsilon^{m^k}$ on m^k has the poles below the real r -axis near $r = 1$. In fact the poles of $dV_\epsilon^{m^k}$ are exactly the same as before, i.e., $d_\epsilon^2 - |x|^2 = 0$: Let $e_1, \dots, e_k, e_{k+1}, \dots, e_n$ be a local orthonormal frame so that $\{e_1, \dots, e_k\}$ spans the tangent space of m^k and $\alpha^1, \dots, \alpha^k, \alpha^{k+1}, \dots, \alpha^n$ be its dual frame so that $AE = I, A = (\alpha_{ij}), E = (e_{ij})$, where $e_j = \sum e_{ij} \frac{\partial}{\partial x_i}$ and $\alpha^i = \sum \alpha_{ij} dx^j$. If we denote the metric tensor g by $G = (g_{ij}), g_{ij} = g(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j})$, then $E^t G E = I$ and hence

$$\begin{aligned}dV^n &= \alpha^1 \wedge \dots \wedge \alpha^n = \det A \quad dx^1 \wedge \dots \wedge dx^n = \frac{1}{\det E} \quad dx^1 \wedge \dots \wedge dx^n \\ &= \sqrt{\det G} \quad dx^1 \wedge \dots \wedge dx^n = \frac{d_\epsilon dx^1 \wedge \dots \wedge dx^n}{(d_\epsilon^2 - |x|^2)^{\frac{n+1}{2}}}.\end{aligned}$$

Since $A = E^{-1} = \frac{1}{\det E} \operatorname{adj} E = \sqrt{\det G} \operatorname{adj} E$, the only poles of $dV^k = \alpha^1 \wedge \cdots \wedge \alpha^k$ are those of dV^n coming from $\sqrt{\det G}$. $\square$

Remark 4.4 *If we consider a geodesic sphere S^k tangent to $\partial\mathbb{H}^n$ and a region U contained in S^k , we can show Theorem 4.3 holds for even more general type region U . In fact, if we let S^k be the k dimensional subspace given by $\{x_1, \dots, x_k, 0, \dots, 0, 1\}$ which is tangent to $\partial\mathbb{H}^n$ at $p = (0, \dots, 0, 1)$ in K^n , then*

$$dV_\epsilon^k = \frac{c_\epsilon dx_1 \wedge \cdots \wedge dx_k}{(c_\epsilon^2 - (x_1^2 + \cdots + x_k^2))^{\frac{k+1}{2}}} = \frac{c_\epsilon r^{k-1}}{(c_\epsilon^2 - r^2)^{\frac{k+1}{2}}} d\theta dr,$$

where $c_\epsilon^2 = d_\epsilon^2 - 1$. If $\epsilon \rightarrow 0$, dV_ϵ^k becomes 0 outside the origin and the origin itself becomes a pole. And it can be shown by direct computation that $\mu_k(U) = 0$ if $p \notin \overline{U}$ and $\mu_k(U) = \frac{1}{2} \operatorname{vol}(\mathbb{S}_H^k)$ if $p \in \operatorname{Int}(U)$. Hence in this case it is clearly invariant under isometry. This singular measure μ_k has some interesting properties, which we do not want to pursue in this paper, but we will state some of these which is related to the theorem. The proofs are direct calculations. Even if μ_k is supported only at the origin, it is not Dirac measure since it is only finitely additive. Indeed if U is a region with piecewise smooth boundary with origin at the boundary point, then $\mu_k(U)$ is determined by the infinitesimal solid angle at the origin since it can be shown easily that μ_k is dilation invariant. This scale invariance also shows that μ_k is invariant under similarity fixing the origin and hence μ_k is invariant under isometry action since the derivatives of the isometries fixing the origin are conformal at the origin.

When we consider a k -dimensional totally geodesic region U , we may assume that U lies on $\mathbb{S}_H^{k+1} = \mathbb{S}_H^n \cap \{x | x_{k+2} = x_{k+3} = \cdots = x_n = 0\}$ by an isometry. Now we have 3 cases as in the 1-dimensional case already discussed: The k -dimensional subspace S^k containing U (i) meets $\partial\mathbb{H}^{k+1}$ transversely, (ii) is tangent to $\partial\mathbb{H}^{k+1}$, or (iii) does not meet $\partial\mathbb{H}^{k+1}$.

For the case (i), we may assume U lies on $\mathbb{S}_H^k$, and for the case (iii), S^k is contained in $\mathbb{L}^{k+1}$ and we may assume U lies on the equator of $\mathbb{S}_H^{k+1}$ which is essentially the same as the standard unit sphere $\mathbb{S}^k$ but with i^k -factor for volume.

The case (ii), the μ_k -measure of total space S^k has value $\operatorname{vol}(\mathbb{S}_H^k)$ from Remark 4.4. So we summarize as follows.

Proposition 4.5 *The k -dimensional μ -measure of any k -dimensional geodesic sphere in $\mathbb{S}_H^n$ is $\operatorname{vol}(\mathbb{S}_H^k)$.*

Remark 4.6 *For a curve γ in $\mathbb{S}_H^n$ especially when it passes through $\partial\mathbb{H}^n$ transversally, it would be natural to define its length by*

$$\mu(\gamma) = \lim_{\epsilon \rightarrow 0} \int_\gamma ds_\epsilon.$$

Of course, we have to show that this definition is invariant under isometry. We do not pursue this issue further in this paper.

Remark 4.7 It is well known that we can use cross ratio to define the distance between the two points (or the length of geodesic line segment) in $\mathbb{H}^2$, and also can extend this to $\mathbb{S}_H^2$ (see [16]). But in this case there arises a subtle and confusing choice problem for multi-valued logarithm.




Fig 4.4

For given two points A and B in $\mathbb{H}^2 \subset K^2$, let P and Q be the points of intersection of $\partial\mathbb{H}^2$ and the line passing through A and B . We assign P and Q so that $\overrightarrow{AB}$ and $\overrightarrow{PQ}$ have the same direction. In this case the hyperbolic distance between A and B is known to be as the following.

$$d_H(A, B) = -\frac{1}{2} \log(PQ|AB) = -\frac{1}{2} \log \frac{\overline{PA}/\overline{AQ}}{\overline{PB}/\overline{BQ}} \quad (7)$$

Here $\overline{PA}, \dots$ etc are Euclidean oriented length so that $\overline{PA} = -\overline{AP}$ and $\overline{xy}$ is positive when $\overrightarrow{xy}$ has the same direction as $\overrightarrow{AB}$.

Now let's use the formula (7) as a definition of distance for any two points in K^2 . Then in general one of the multi-values of $\log$ in (7) coincide with μ -distance, but it is unclear which choices are natural and consistent for all cases.

Let $A = (x, 0)$ and $B = (\frac{1}{x}, 0)$. Then $d_H(A, B) = -\frac{1}{2} \log(-1) = \pm \frac{\pi}{2}i + m\pi i$, $m \in \mathbb{Z}$. Taking the clockwise contour from 1 to -1 , gives the value $\frac{\pi}{2}i$, which coincides with μ -distance of A and B .

If the line passing through A and B is tangent to $\partial\mathbb{H}^2$ at $P = Q$, we have to carry out a formal computation for (7) and choose $\log$ as follows to obtain μ -distance; $\log \frac{0/0}{1/-1} = \log(-1) = -\pi i$ or $\log 1 = -2\pi i$ depending on whether $A = P = Q \neq B$ or $A \neq P = Q \neq B$ ($A < P < B$), respectively.

The remaining case is when the line extending AB does not meet $\partial\mathbb{H}^2$. Let the equation of this line be $x_2 = k$ ($k > 1$) and $A = (a, k), B = (b, k)$ with $a < b$ on K^2 . If we solve simultaneously with $x_1^2 + x_2^2 = 1$, we have $x_1 = \pm\sqrt{k^2 - 1}i$. Now from (7) with $P = (\sqrt{k^2 - 1}i, k)$ and $Q = (-\sqrt{k^2 - 1}i, k)$ or $P = (-\sqrt{k^2 - 1}i, k)$ and $Q = (\sqrt{k^2 - 1}i, k)$, we get $d_H(A, B) = \pm\alpha i + m\pi i$ where $m \in \mathbb{Z}$, $0 < \alpha < \pi$. Among these, the choice αi coincides with the μ -distance and we should have chosen $P = (\sqrt{k^2 - 1}i, k)$ and $Q = (-\sqrt{k^2 - 1}i, k)$ to

obtain αi , and it looks confusing to explain why this is a consistent choice. The other choice of P and Q corresponds to the choice $\tilde{d}_\epsilon = 1 + \epsilon i$ for ϵ -approximation of our singular metric.

The extended Kleinian model $\mathbb{H}_K^n$ has a projective geometric structure, so a geodesic in $\mathbb{H}_K^n$ is a straight line and a dual of a point x , i.e., $x^\perp$ is easily obtained as usual (see Fig 4.5 and 4.6). Then the length of a geodesic line segment joining x (respectively y) and an arbitrary point in $x^\perp$ (respectively $y^\perp$) is $\frac{\pi}{2}i$. Because there is an isometry which takes x and $x^\perp$ to a point on the equator and to a longitude respectively, and takes y and $y^\perp$ to a north pole and to the equator respectively.




Fig 4.5




Fig 4.6

Now we define angles on the extended model $\mathbb{S}_H^n$. In fact we can define a notion of angle on a semi-Riemannian manifold using μ -measure on $\mathbb{S}_H^1$. For two tangent vectors v_p and w_p at a point p on a Riemannian manifold, the notion of the angle θ between these two vectors is obvious from that of standard Euclidean plane $\mathbb{R}^2$ and can be calculated by the equation,

$$\langle v_p, w_p \rangle = \|v_p\| \|w_p\| \cos \theta, \quad 0 \leq \theta < \pi. \quad (8)$$

But for a semi-Riemannian manifold, we have some difficulties with this formula since the function $\cos^{-1}$ is multi-valued and θ can take several complex values. The definitions of angle have been given through the combinatorial way in [6] and the cross ratio in [16]. Now that we have a notion of arc length on $\mathbb{S}_H^1$, as μ -measure, we can define an angle just as for Riemannian case.

Definition 4.8 For given two vectors $v, w \in \mathbb{R}^{n,1}$, the angle between v and w , $\theta = \angle(v, w)$, is defined as $-i \cdot d_H(v, w)$, where $d_H(v, w)$ is the length of a geodesic segment joining two points of $\mathbb{S}_H^n$ radially projected from v, w to $\mathbb{S}_H^n$.

Fig. 4.7 shows the various angles of between vectors in a degenerate space which occurs as a tangent space to the light cone. (Two poles in the picture represent light-like vectors of $\mathbb{R}^{n,1}$.)




Fig 4.7

Notice that the factor $-i$ is multiplied to normalize the total length $2\pi i$ of the great circle in $\mathbb{S}_H^n$ as 2π (see Theorem 4.2). If we used $\tilde{d}_\epsilon = 1 + \epsilon i$ for our approximation, we have to use $i d_H(v, w)$ instead of $-i d_H(v, w)$ since the length of great circle in this case becomes $-2\pi i$.

To check the equation (8) for Lorentzian case, we need to define the norm $\|\cdot\|$ first. The Lorentzian norm of a vector v in $\mathbb{R}^{n,1}$ is defined to be a complex number

$$\|v\| = \langle v, v \rangle^{\frac{1}{2}},$$

where $\|v\|$ is either positive, zero, or positive imaginary.

From our definitions of angle and norm, it can be shown that (8) holds for all cases.

Proposition 4.9 *For non-null vectors v and w in the space $\mathbb{R}^{n,1}$, we have*

$$\langle v, w \rangle = \|v\| \|w\| \cos \angle(v, w). \quad (9)$$

Proof It is not hard to check (9) for the various cases through simple computations, we can see the details at [4]. $\square$

Our definition of an angle clearly satisfies the following four properties by Theorem 4.3 and Remark 4.4,

- (i) the invariance under isometry,
- (ii) equation (9),
- (iii) finite additivity of angle: if θ consists of two parts θ_1 and θ_2 , then $\theta = \theta_1 + \theta_2$,
- (iv) the angle of half rotation is π , i.e., a straight line has angle π .

Conversely, it can be shown that the angle is uniquely determined by these four properties.

There are various formulas similar to (9) relating Lorentzian inner product and angle or hyperbolic distance between two points or its dual hyperplane depending on the position of vectors in [18, Proposition 2.4.5], and all of these are equivalent to (9) one single formula.

More generally if we consider the angle between two vectors v and w on $\mathbb{R}^{p,q}$ or two tangent vectors on $\mathbb{S}_H^n$ or on a semi-Riemannian manifold, we have to look at the plane $P = \text{span}\{v, w\}$ spanned by these two vectors.




Fig 4.8

The plane P is isometric to $\mathbb{R}^{2,0}$, $\mathbb{R}^{1,1}$, or $\mathbb{R}^{0,2}$ for non-degenerate cases and to $\mathbb{R}^{1,0}$, $\mathbb{R}^{0,1}$, or $\mathbb{R}^{0,0}$ for degenerate cases depending on the situations. At a point p on $\mathbb{S}_H^n$ not lying on $\partial\mathbb{H}^n$, the angle between the two tangent vectors v_p and w_p is determined on the plane $\text{span}\{v'_p, w'_p\}$ where p' is the radial projection of p on the hyperboloid $H_\pm^n$ or S_1^n and $v'_p, w'_p \in T_{p'}(H_\pm^n)$ or $T_{p'}(S_1^n)$ are projections of v_p, w_p respectively. The plane $\text{span}\{v'_p, w'_p\}$ has signature $\mathbb{R}^{2,0}, \mathbb{R}^{1,1}$ or $\mathbb{R}^{1,0}$ depending on the position of p , and in each case the angle is determined in the usual way for the definite cases and using Definition 4.8 for $\mathbb{R}^{1,1}$ and $\mathbb{R}^{1,0}$ cases.

If $p \in \mathbb{S}_H^n$ lies in the Lorentzian part S_1^n , the norm $\|v_p\|$ of a tangent vector v_p and the norm of its translate to the origin, denoted by v' , are related by, $\|v'\| = -i\|v_p\|$ from our sign conventions (see Fig. 4.8). And $\langle v', w' \rangle = -\langle v_p, w_p \rangle$ by the definition of the metric on S_1^n . Hence we have the following identity

$$\frac{\langle v_p, w_p \rangle}{\|v_p\| \|w_p\|} = \frac{\langle v', w' \rangle}{\|v'\| \|w'\|}, \quad (10)$$

and $\angle(v_p, w_p) = \angle(v', w')$ by the four characterizing properties of angle.

Proposition 4.10 *For $p \in \mathbb{S}_H^n$ and $p \notin \partial\mathbb{H}^n$, the angle $\angle(v_p, w_p)$ between two tangent vectors $v_p, w_p \in T_p\mathbb{S}_H^n$ is equal to $-id_H(v', w')$, where v' is a vector at the origin of $\mathbb{R}^{n,1}$ given as a parallel translation of the tangent vector of $T_p H_\pm^n$ or $T_p S_1^n$ which is obtained as a radial projection of v_p . (We also can see the point of K^n determined by v' as in Fig. 4.9.)*

Proof If p is in the Lorentzian part, we may assume p lies on the equator via isometry. In this case, $v' = v$ and $w' = w$, and $\angle(v_p, w_p) = \angle(v, w) = -i d_H(v, w) = -i d_H(v', w')$ hold from (10) and Definition 4.8.

If p is in the hyperbolic part, we may assume p is the north pole and we clearly have $d_H(v', w') = d_H(v, w) = i\angle(v_p, w_p)$. $\square$




Fig 4.9

The isometry invariance of angle at a point $p \in \mathbb{S}_H^n \setminus \partial\mathbb{H}^n$ is obtained from Proposition 4.10.

If $p \in \mathbb{S}_H^2$ lies on $\partial\mathbb{H}^2$, we define angle as follows.

Definition 4.11 *We define the angle $\angle(v_i, v_{i+1})$ for the tangent vectors $v_1, \dots, v_5$ configured as in Fig. 4.10:*

$$\angle(v_1, v_2) = \angle(v_5, v_1) = \frac{\pi}{2}, \quad \angle(v_2, v_3) = \angle(v_4, v_5) = 0, \quad \text{and} \quad \angle(v_3, v_4) = \pi.$$




Fig 4.10

Remark 4.12 *A justification of this definition can be given as follows. For $p \in \partial\mathbb{H}^2$ in K^2 , we first consider θ_ϵ from the equation (8) using ϵ -metric and then determine θ as $\lim_{\epsilon \rightarrow 0} \theta_\epsilon$. In this process we again have a problem of choice of values for multi-valued function $\cos^{-1}$. But our choice is the simplest reasonable one satisfying the four characterizing of angle.*

If $p \in \partial\mathbb{H}^n$ with $n \geq 3$, we have another type of tangent plane at p which touches $\partial\mathbb{H}^n$ at the only point p . In this case, the ϵ -metric and hence the limit has a rotational symmetry around p and hence the angle is the same as the usual Euclidean angle.

Remark 4.13 When we consider an angle on $\mathbb{S}_H^n$, we have 3 different approaches other than using (Definition 4.8, 4.11) or (Proposition 4.10, Definition 4.11) or (four characterizing properties of angle).

One way is to use the dual distance and we will defer the discussion of this topic to elsewhere.

Secondly we can define the angle between v_p and w_p as the area of the lune determined by these two vectors. The angle is given by $-\frac{1}{2} \times$ the area (2-dimensional μ -measure) of the lune.

Third definition uses cross ratio and will be explained separately in the next remark.

Remark 4.14 As we explained distance on K^2 using cross ratio, we also can explain angle using cross ratio. Let P be a point in the Lorentzian part, and θ be an angle between two lines a and b . Let s and t be two lines through p tangent to $\partial\mathbb{H}^2$, and be ordered so that the directions $\overrightarrow{ab}$ and $\overrightarrow{st}$ match.




Fig 4.11

Choose any two lines $l_i, i = 1, 2$ and let S_i, A_i, B_i, T_i be the intersecting points of l_i and the lines s, a, b, t respectively. Then it is well known from a Pappus theorem that the cross ratio $(S_i T_i | A_i B_i)$ agrees for $i = 1, 2$, and hence we can define $(st|ab) := (S_i T_i | A_i B_i)$ and the angle θ is defined as $\angle\theta = \frac{i}{2} \log(st|ab)$ (see [16]).

Now it can be shown using Remark 4.7 and Proposition 4.10 that this definition agrees with ours.

If p lies inside $\mathbb{H}^2$, we can compute the tangent lines s and t through complexification. In this case, we have some ambiguity for the choice of s and t as in the later half of Remark 4.7. Anyway Remark 4.7 and Proposition 4.10 also imply that our angle θ agrees with an appropriate choice of angle coming from the cross ratio.

When p lies on $\partial\mathbb{H}^2$, the two tangent lines s and t coincide and we can derive the same value as in the Fig. 4.10 from the cross ratio by choosing an appropriate values of $\log$.

5 Gauss volume formula and Santaló-Milnor-Suárez-Peiró relation in the extended hyperbolic model

Choose a point $x \in \mathbb{S}_H^n$ and consider its dual hypersphere $x^\perp \cap \mathbb{S}_H^n$. For a subset $B \subset x^\perp \cap \mathbb{S}_H^n$, the union of the great semi-circles passing through $b \in B$ connecting x and $-x$ will be called a lune $L = L(x, B)$ with vertex x and cross section B . And the $(n-1)$ -dimensional μ -measure $\mu(B)$ in $x^\perp \cap \mathbb{S}_H^n$ will be called a hyperbolic solid angle of L . In fact, if $x = e_0 = (1, 0, \dots) \in \mathbb{R}^{n,1}$, then the dual hypersphere is $\mathbb{S}_1^n \cap \{x \in \mathbb{R}^{n,1} | x_0 = 0\}$ and the radial projection (or an inverse image of exponential map) of a lune $L(e_0, B)$ becomes a cone in the tangent space $T_{e_0}(\mathbb{S}_H^n)$ and $\mu(B)$ is i^{n-1} times the usual solid angle of the cone in Euclidean space $T_{e_0}(\mathbb{S}_H^n)$, i.e., i^{n-1} times spherical volume of the portion in $\mathbb{S}^{n-1}$ cut out by the cone. We should be careful when we consider the volume of $\mathbb{S}^{n-1}$ in $\mathbb{S}_H^n$ since it lies in $\mathbb{L}^n$ and its metric is the negative of the usual one and the volume is determined as $i^{n-1} \cdot \text{vol}(\mathbb{S}^{n-1})$ by Convention and of course is equal to $\text{vol}(\mathbb{S}_H^{n-1})$.

Theorem 5.1 *A lune of hyperbolic solid angle S in $\mathbb{S}_H^n$ has volume $\frac{\text{vol}(\mathbb{S}_H^n)}{\text{vol}(\mathbb{S}_H^{n-1})} \cdot S$, where the vertex of the lune assumed not to lie on $\partial\mathbb{H}^n$ for convenience. (See the following remark for the case when the vertex lies on $\partial\mathbb{H}^n$.)*

Proof Let's consider first the case when the vertex of a lune lies in $\mathbb{H}^n$. In this case we may assume the vertex is $e_0 = (1, 0, \dots)$, i.e., vertex is $(0, \dots, 0) \in K^n$, via isometry, and the volume of a lune is clearly proportional to the solid angle by the rotational symmetry of $\mathbb{S}_H^n$.

When the vertex of a lune lies in the Lorentzian part S_1^n , we may assume the vertex is $e_1 = (0, 1, 0, \dots)$, i.e., vertex is $(\infty, 0, \dots, 0) \in K^n$, by an isometry. Then such lune L can be given as a cylinder of type $(-\infty, \infty) \times S$ in the Kleinian model, and the volume of L will be given by

$$\mu(L) = \lim_{\epsilon \rightarrow 0} \int_S \int_{-\infty}^{\infty} \frac{d_\epsilon dx_1}{(d_\epsilon^2 - x_1^2 - x_2^2 - \dots - x_n^2)^{\frac{n+1}{2}}} dx_2 \cdots dx_n \quad (n \geq 2).$$

To compute the integral, we divide S into two parts A and B depending on whether $x_1^2 + x_2^2 + \dots + x_n^2$ is greater than 1 and less than or equal to 1 respectively. Since the sign of the volume of Lorentzian part is i^{n+1} , the integral $\int_{-\infty}^{\infty} \frac{d_\epsilon dx_1}{(d_\epsilon^2 - x_1^2 - x_2^2 - \dots - x_n^2)^{\frac{n+1}{2}}}$ in the above formula for A part can be replaced by $i^{n+1} \int_{-\infty}^{\infty} \frac{d_\epsilon dx_1}{(x_1^2 + x_2^2 + \dots + x_n^2 - d_\epsilon^2)^{\frac{n+1}{2}}}$ when the integrand has positive real part (and positive as $\epsilon \rightarrow 0$). If we let $a = x_2^2 + \dots + x_n^2 - d_\epsilon^2$ and $x_1 = a^{\frac{1}{2}}t$ with

$Re(a^{\frac{1}{2}}) > 0$, then

$$\begin{aligned} i^{n+1} \int_{-\infty}^{\infty} \frac{dx_1}{(x_1^2 + a)^{\frac{n+1}{2}}} &= \frac{i^{n+1}}{a^{\frac{n}{2}}} \int_{\gamma_1} \frac{2 dt}{(1+t^2)^{\frac{n+1}{2}}} = \frac{2i^{n+1}}{a^{\frac{n}{2}}} \int_0^{\infty} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} \\ &= \frac{2i}{(d_{\epsilon}^2 - x_2^2 - \dots - x_n^2)^{\frac{n}{2}}} \int_0^{\infty} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}}. \end{aligned}$$

Here $\frac{1}{(d_{\epsilon}^2 - x_2^2 - \dots - x_n^2)^{\frac{n}{2}}} = \frac{i^n}{a^{\frac{n}{2}}}$. Since A is a Lorentzian part in K^{n-1} and γ_1 is a contour given in Fig. 5.1.

For the B part, let $b = d_{\epsilon}^2 - x_2^2 - \dots - x_n^2$ and $x_1 = b^{\frac{1}{2}}y$ with $Re(b^{\frac{1}{2}}) > 0$, then

$$\int_{-\infty}^{\infty} \frac{dx_1}{(b - x_1^2)^{\frac{n+1}{2}}} = \frac{2}{b^{\frac{n}{2}}} \int_{\gamma_2} \frac{dy}{(1-y^2)^{\frac{n+1}{2}}} = \frac{2}{b^{\frac{n}{2}}} \int_{\gamma_3} \frac{dy}{(1-y^2)^{\frac{n+1}{2}}}.$$

Substituting $y = it$, we obtain

$$\frac{2}{b^{\frac{n}{2}}} \int_{\gamma_3} \frac{dy}{(1-y^2)^{\frac{n+1}{2}}} = \frac{2i}{b^{\frac{n}{2}}} \int_{\gamma_4} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} = \frac{2i}{b^{\frac{n}{2}}} \int_0^{\infty} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}}.$$




Fig 5.1

Claim. $\int_0^{\infty} \frac{dt}{(1+t^2)^{\frac{n+1}{2}}} = \frac{\text{vol}(\mathbb{S}^n)}{2\text{vol}(\mathbb{S}^{n-1})} \quad (n \geq 2).$

Now if assume this claim, we can write for both A and B parts as

$$\mu_n(L) = \lim_{\epsilon \rightarrow 0} \frac{i \text{vol}(\mathbb{S}^n)}{\text{vol}(\mathbb{S}^{n-1})} \int_S \frac{d_{\epsilon} dx_2 \cdots dx_n}{(d_{\epsilon}^2 - x_2^2 - \dots - x_n^2)^{\frac{n}{2}}} = \frac{\text{vol}(\mathbb{S}_H^n)}{\text{vol}(\mathbb{S}_H^{n-1})} \mu_{n-1}(S),$$

where $\mu_n(L)$ is the n -dimensional volume or μ -measure. And this completes the proof of the theorem.

Proof of Claim Let $I_m = \int_0^{\infty} \frac{dt}{(1+t^2)^m}$. Then $I_m = \frac{2m-3}{2m-2} I_{m-1}$ ($m \geq 2$) is induced from

$$\int \frac{dt}{(1+t^2)^m} = \frac{1}{2(m-1)} \frac{t}{(1+t^2)^{m-1}} + \frac{2m-3}{2m-2} \int \frac{dt}{(1+t^2)^{m-1}}.$$

Prove the Claim using induction on $n \geq 2$. If $n = 2$, $I_{3/2} = [\frac{t}{\sqrt{1+t^2}}]_0^\infty = 1 = \frac{\text{vol}(\mathbb{S}^2)}{2\text{vol}(\mathbb{S}^1)}$.

And if $n = 3$, $I_2 = \frac{1}{2}I_1 = \frac{\pi}{4} = \frac{\text{vol}(\mathbb{S}^3)}{2\text{vol}(\mathbb{S}^2)}$. Now using the hypothesis, we have

$$I_{\frac{n+2}{2}} = \frac{n-1}{n} I_{\frac{n}{2}} = \frac{n-1}{n} \frac{\text{vol}(\mathbb{S}^{n-1})}{2\text{vol}(\mathbb{S}^{n-2})} = \frac{\text{vol}(\mathbb{S}^{n+1})}{2\text{vol}(\mathbb{S}^n)}.$$

The last equality follows from the well-known formula

$$\frac{\text{vol}(\mathbb{S}^n)}{\text{vol}(\mathbb{S}^{n-2})} = \frac{2\pi}{n-1} \quad \text{or} \quad \text{vol}(\mathbb{S}^n) = \frac{2\pi^{\frac{n+1}{2}}}{\Gamma(\frac{n+1}{2})}.$$

□

Remark 5.2 *The above theorem seems still hold even when the vertex of a lune lies on the ideal boundary if the lune is transversal to the ideal boundary. The non-transversal case looks rather complicated and subtle. We illustrate these when the dimension $n = 2$ to show some ideas.*




Fig 5.2

The first picture of Figure 5.2 is a lune L with its vertex $x \in \partial\mathbb{H}^2$ and its angle 0. We can change L into a region consisting of a lune L_0 with its vertex at the origin and angle α and two right triangles as in the second picture so that it clearly has the same area as L . Now from this, we can show the area of L is in fact zero as follows.

$$\begin{aligned} \mu(L) &= \lim_{\epsilon \rightarrow 0} \int_L dV_\epsilon \\ &= \lim_{\epsilon \rightarrow 0} \int_{\triangle OCD \cup \triangle OC'D' \cup L_0} dV_\epsilon \\ &= 2\mu(\triangle OCD) + \mu(L_0) \\ &= 2(\pi - \frac{\pi}{2} - (\frac{\pi}{2} - \alpha) - 0) + (\frac{-4\pi}{2\pi i})\alpha i \\ &= 0. \end{aligned}$$




Fig 5.3

On the other hand, for this L , $x^\perp \cap L = \{x\}$ and hence $\mu(B) = 0$. Therefore we see that the theorem holds for such L .

If a Lune L has an angle π as in the figure 5.3 (i), then its complement $K^2 \setminus L$ is a lune of angle 0. Hence we have

$$\mu(L) = -2\pi = \frac{-4\pi}{2\pi i} \cdot \pi i = \frac{\text{vol}(\mathbb{S}_H^2)}{\text{vol}(\mathbb{S}_H^1)} \mu_1(x^\perp \cap L)$$

and the theorem holds.

If a lune L has an angle $\frac{\pi}{2}$ as in figure 5.3 (ii), then let S_3 and S_4 are the reflection of S_2 and S_1 respectively about the vertical line through x . Even if $\mu(S_1)$ and $\mu(S_2)$ are not defined separately, we can compute $\mu(L)$ as follows using the existence of $\mu(S_1 \cup S_3)$.

$$\begin{aligned} \mu(L) &= \lim_{\epsilon \rightarrow 0} \int_{S_1 \cup S_2} dV_\epsilon = \lim_{\epsilon \rightarrow 0} \left(\int_{S_1} dV_\epsilon + \int_{S_2} dV_\epsilon \right) \\ &= \lim_{\epsilon \rightarrow 0} \left(\int_{S_1} dV_\epsilon + \int_{S_3} dV_\epsilon \right) = \lim_{\epsilon \rightarrow 0} \int_{S_1 \cup S_3} dV_\epsilon \\ &= \mu(S_1 \cup S_3) = (\mu(S_1 \cup S_3) + \mu(S_2 \cup S_4))/2 \\ &= -\pi. \end{aligned}$$

In this case, it is unclear how to define a solid angle. But the above result suggests that it would be reasonable to interpret that L occupies the half of the total solid angle.

Remark 5.3 When the dimension $n = 2$, we have to be careful about the notions of angle and solid angle. If the angle of a lune is θ , then its solid angle becomes θi from its definition.

When the dimension n is even, the volume of a spherical simplex on $\mathbb{S}^n$ can be obtained combinatorially as an alternating sum of its solid angle using the well known Euler-Poincaré method (see [1, p.120]). And then it can be shown that the same formula (only differ by

sign) holds for a hyperbolic simplex indirectly using an analytic continuation technique. But if we use the extended model $\mathbb{S}_H^n$, the above Euler-Poincaré method can be applied directly without any change showing that the above formula holds not only for a hyperbolic simplex but also for a Lorentzian simplex or even for a simplex lying across the ideal boundary $\partial\mathbb{H}^n$, that is, any simplex on $\mathbb{S}_H^n$.

When we consider the formula, it is more convenient to use the normalized volume so that the total volume becomes 1. The normalized volume of a lune L is given by $\widehat{\mu}(L) = \frac{\mu(L)}{\text{vol}(\mathbb{S}_H^n)}$ and the normalized solid angle of L is given by $\widehat{\alpha}(L) = \frac{\alpha(L)}{\text{vol}(\mathbb{S}_H^{n-1})}$.

Given an n -simplex Δ^n on $\mathbb{S}_H^n$, Δ^n is an intersection of half spaces $H_0, H_1, \dots, H_n$ and $H_{i_1} \cap \dots \cap H_{i_k}$ ($k \leq n$) becomes a lune $L_{i_1 \dots i_k}$. Then the normalized volume of Δ^n can be given as follows using Euler-Poincaré method and Theorem 5.1.

$$\begin{aligned} \widehat{\mu}(\Delta^n) &= \frac{1}{2} \sum_{k=0}^n \sum_{i_1 < \dots < i_k} (-1)^k \widehat{\mu}(L_{i_1 \dots i_k}) \\ &= \frac{1}{2} \sum_{k=0}^n \sum_{i_1 < \dots < i_k} (-1)^k \widehat{\alpha}(L_{i_1 \dots i_k}) \\ &=: \frac{1}{2} k(\Delta^n). \end{aligned} \tag{11}$$

If we apply the formula (11) when $n = 2$ for a triangle Δ on $\mathbb{S}_H^2$ with three angles A, B , and C , then we have

$$\frac{\mu(\Delta)}{-4\pi} = \widehat{\mu}(\Delta) = \frac{1}{2} \left(1 - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} + \frac{Ai}{2\pi i} + \frac{Bi}{2\pi i} + \frac{Ci}{2\pi i} \right).$$

It follows that $\mu(\Delta) = \pi - A - B - C$, and we see that this beautiful formula can be extended even across the ideal boundary $\partial\mathbb{H}^2$. This was first obtained by J. Böhm and Im Hof [2].

Suppose M is a hyperbolic manifold with a triangulation $\mathcal{T}$ consisting of totally geodesic n -simplex. Then it can be shown easily that $\chi(M) = \sum_{\Delta^n \in \mathcal{T}} k(\Delta^n)$ (see [7]). Therefore the Gauss-Bonnet theorem for a hyperbolic manifold follows immediately.

$$\begin{aligned} \mu(M^n) &= \text{vol}(M^n) = \text{vol}(\mathbb{S}_H^n) \sum_{\Delta^n \in \mathcal{T}} \widehat{\text{vol}}(\Delta^n) = \text{vol}(\mathbb{S}_H^n) \sum_{\Delta^n \in \mathcal{T}} \widehat{\mu}(\Delta^n) \\ &= \frac{\text{vol}(\mathbb{S}_H^n)}{2} \sum_{\Delta^n \in \mathcal{T}} k(\Delta^n) = \frac{\text{vol}(\mathbb{S}_H^n)}{2} \chi(M^n). \end{aligned} \tag{12}$$

Therefore we conclude the following result from the Euler-Poincaré method used in $\mathbb{S}_H^n$.

Proposition 5.4 *When the dimension n is even, we have*

$\widehat{\mu}(\Delta^n) = \frac{1}{2} \sum_{k=0}^n \sum_{i_1 < \dots < i_k} (-1)^k \widehat{\alpha}(L_{i_1 \dots i_k})$ for an n -simplex Δ^n in $\mathbb{S}_H^n$ and $\mu(M^n) = \frac{\text{vol}(\mathbb{S}_H^n)}{2} \chi(M^n)$ for a hyperbolic n -manifold. In particular, the area of a triangle in $\mathbb{S}_H^2$ is $\pi - A - B - C$, where A, B, C are the angles of the given triangle.

Similarly if we consider a Lorentzian spherical manifold M (i.e., metric signature $(-, +, \dots, +)$ with constant sectional curvature $\mathcal{K} \equiv 1$, or metric $(+, -, \dots, -)$ with $\mathcal{K} \equiv -1$), then M has a developing on the Lorentzian part and we have the same formula $\mu(M^n) = \frac{\text{vol}(\mathbb{S}_H^n)}{2} \chi(M^n)$.

In this argument we have to be careful not to have a simplex Δ^n whose (extended) face is tangent to $\partial\mathbb{H}^n$ since the solid angles are not defined for such a simplex. But of course this can be easily achieved by perturbing the triangulation.

Now notice that $\mu(M^n)$ has sign $-i^{n-1} = i^{n+1}$ and $\text{vol}(\mathbb{S}_H^n)$ has sign i^n . This shows that we have both $\mu(M^n) = 0$ and $\chi(M^n) = 0$. We already know that latter should hold since M is Lorentzian, but the condition $\mu(M^n) = 0$ is absurd and we can conclude that there does not exist such closed manifold M . Of course, this fact has been known and can be deduced from the usual Gauss-Bonnet theorem for semi-Riemannian manifolds (see [9] for instance), but we could see this immediately by an elementary combinatorial way.

Proposition 5.4 gives us an interesting consequences especially when the $\Delta(A, B, C)$ lies across the ideal boundary $\partial\mathbb{H}^2$ such that the edges are not completely contained in the Lorentzian part. In this case $\pi - A - B - C$ is a complex number whose real part is the area of truncated polygon and whose imaginary part is the length of the edge introduced in the truncation as the following examples show.

Example 5.5 *The area of the triangles in Fig. 5.4 and 5.5 are $a \cdot i = \pi - (\frac{\pi}{2} + \frac{\pi}{2} - ia)$ and $-A = \pi - (\frac{\pi}{2} + \frac{\pi}{2} + A)$ respectively.*




Fig 5.4




Fig 5.5

Example 5.6 *The triangle $(1,2,3)$ in Fig 5.6 has area $\pi - (A - bi - ei) = \pi - A + (b + e)i$. Also the triangle $(1,2,3)$ is divided into three polygons $(2,4,5)$, $(3,6,7)$ and $(1,4,5,7,6)$ and each polygon has pure imaginary area ei , bi and real value area $\pi - A$ respectively.*




Fig 5.6




Fig 5.7

Example 5.7 The triangle $(1,2,3)$ in Fig 5.7 is divided into various polygons. Notice that the line segment $(6,7)$ is $3^\perp$ and $(4,5)$ is $2^\perp$, and hence $(2,3)$ is $8^\perp$ which implies that the length of the line segment $(5,8)$ is equal to $\frac{\pi}{2}i$ and similarly for $(7,8)$. Let b be the length of $(4,8)$, c for $(6,8)$ and α be the angle $\angle(4,8,6)$. Then we have

$$\begin{aligned} \text{Area } (1, 2, 3) &= \pi - (A + (\frac{\pi}{2} - bi) + (\frac{\pi}{2} - ci)) \\ &= -A + (b + c)i, \end{aligned}$$

and

$$\begin{aligned} &\text{Area } (1, 4, 8, 6) + \text{Area } (2, 8, 3) + \text{Area } (2, 4, 8) + \text{Area } (3, 6, 8) \\ &= (\pi - A - \alpha) + (\alpha - \pi) + (bi) + (ci) \\ &= -A + (b + c)i \end{aligned}$$

Notice that the real part is the area of the shaded quadrangle and the imaginary part is the total length of the truncated sides of the quadrangle.

Example 5.8 From the outer triangle which is dual to the inner triangle $\triangle(1,2,3)$ in Fig 5.8, we get

$$\text{Area } (1', 2', 3') = \pi - (\pi - ai + \pi - bi + \pi - ci) = -2\pi + (a + b + c)i.$$

If the inner triangle in the figure shrinks to one point, then the area of the triangle becomes $-2\pi (= \text{vol } (\mathbb{S}_H^2)/2)$. In fact in this case, the vertices of the triangle lies on the equator and the angle at the vertices become π .

We can generalize hyperbolic trigonometry to any triangle in the extended hyperbolic space $\mathbb{S}_H^2$ or in general $\mathbb{S}_H^n$ (see [4]). Naturally, already known cosine and sine laws at the hyperbolic space can be applicable to the extended hyperbolic space. Then all the known trigonometric formulas for special hyperbolic polygons are in fact explained in a consistent manner through the trigonometry for various triangles of $\mathbb{S}_H^2$ lying across the ideal boundary.




Fig 5.8

See Fig. 5.6 and 5.7 for the right angled polygons and the corresponding triangles in $\mathbb{S}_H^2$. All the cosine and sine laws of quadrangles in Fig. 5.6 and 5.7 could be obtained from the cosine and sine laws of the outer triangles $\triangle(1, 2, 3)$ in the figures. For example, the following cosine law of the pentagon in Fig. 5.6 $\cosh a = \frac{\cosh b \cosh e + \cos A}{\sinh b \sinh e}$ is deduced from the following generalized cosine law for the triangle $\triangle(1, 2, 3)$ in $\mathbb{S}_H^2$,

$$\cosh(a + \pi i) = \frac{\cos(-bi) \cos(-ei) + \cos A}{\sin(-bi) \sin(-ei)}.$$

Also the extended hyperbolic space shows an evident and geometric reason of resemblance of laws between the hyperbolic space and spherical space. Furthermore we can as well derive the trigonometry for the Lorentzian triangles and polygons, which has not been determined well as far as we know. We refer the interested reader to see [4] for the details.

From Theorem 5.1, we can also induce the volume of elementary three dimensional objects, such as lens and trihedron $\text{Tri}(A, B, C)$ with dihedral angles A, B, C .

Corollary 5.9 *The 3-dimensional volume of a convex lens $l(A)$ with dihedral angle A is $-A\pi i$.*

Proof

$$\begin{aligned} & \frac{\text{vol}(\mathbb{S}_H^3)}{\text{vol}(\mathbb{S}_H^2)} \times \text{Area of 2-dimensional cross section for the lens of angle } A \\ &= \frac{\pi}{2} i \times (-2A). \end{aligned}$$

□

Corollary 5.10 *The volume of a trihedron $\text{Tri}(A, B, C)$ with dihedral angles A, B, C is represented as*

$$\text{vol}(\text{Tri}(A, B, C)) = \frac{\pi}{2}i(\pi - A - B - C).$$

The volume of a hyperbolic trihedron has a similar expression as the spherical one which has a well-known volume formula, $V = \frac{\pi}{2}(A + B + C - \pi)$. The difference between the two formulas is i^3 .

Example 5.11 *Consider a tetrahedron Δ^3 some of whose vertices (but not edges) are lying outside the hyperbolic part. Let P be a hyperbolic polyhedron truncated from Δ^3 by the dual planes of outside vertices and T be the truncated part of Δ^3 so that $\Delta^3 = P \cup T$. See Fig. 5.9. Then we have*

$$\begin{aligned} \text{vol}(\Delta^3) &= \text{Re vol}(\Delta^3) + \text{Im vol}(\Delta^3) \\ &= \text{vol}(P) + \text{vol}(T) \\ &= \text{vol}(P) + \frac{\pi}{4}i \times (\text{area of the truncated face}). \end{aligned}$$

This follows immediately from Theorem 5.1 since T is exactly the half of the lune containing T .




Fig 5.9

Ushijima [19] found that $\text{vol}(P)$ is the real part of the value which is obtained by applying Murakami-Yano's volume formula for hyperbolic tetrahedron [12] formally to Δ^3 . In this case we have a problem of determining a "right" value among the multi-values from the formula and in our extended model the above $\text{vol}(\Delta^3)$ is the correct value and the pure imaginary part is interpreted as the volume of T as well as the area of the truncated face.

The general statement for such phenomenon requires a proof of the analyticity of the volume formula for tetrahedra across the boundary in the extended model and will be deferred to a subsequent paper.

We consider another interesting application of the extended model. Let $\triangle(1, 2, 3)$ be a convex spherical triangle with vertices 1, 2, 3 on the 2-sphere $\mathbb{S}^2 \subset \mathbb{R}^3$ and $\triangle(1', 2', 3')$ be its dual triangle so that the distance between a vertex i' (i , resp.) and each point on the edge (j, k) ((j', k') , resp.) is $\frac{\pi}{2}$, i.e., they are perpendicular as vectors in $\mathbb{R}^3$, where $\{i, j, k\} = \{1, 2, 3\}$. Then $\mathbb{S}^2$ can be decomposed into 8 triangles, $\triangle(1, 2, 3)$, $\triangle(1', 2, 3)$, $\triangle(1, 2', 3)$, $\triangle(1, 2, 3')$, $\triangle(1', 2', 3)$, $\triangle(1', 2, 3')$, $\triangle(1, 2', 3')$, $\triangle(1', 2', 3')$. If we denote the volume of $\triangle(i, j, k)$ again by $\triangle(i, j, k)$ abusing the notation, then we have

$$\begin{aligned} & \triangle(1, 2, 3) + \triangle(1', 2', 3) + \triangle(1', 2, 3') + \triangle(1, 2', 3') \\ &= (A + B + C - \pi) + (\pi - C + \frac{\pi}{2} + \frac{\pi}{2} - \pi) + (\pi - B + \frac{\pi}{2} + \frac{\pi}{2} - \pi) + (\pi - A + \frac{\pi}{2} + \frac{\pi}{2} - \pi) \\ &= 2\pi, \end{aligned}$$

$$\begin{aligned} & \triangle(1', 2, 3) + \triangle(1, 2', 3) + \triangle(1, 2, 3') + \triangle(1', 2', 3') \\ &= (a + \frac{\pi}{2} + \frac{\pi}{2} - \pi) + (b + \frac{\pi}{2} + \frac{\pi}{2} - \pi) + (c + \frac{\pi}{2} + \frac{\pi}{2} - \pi) + (\pi - a + \pi - b + \pi - c - \pi) \\ &= 2\pi, \end{aligned}$$

where A, B, C are the angles at vertices 1, 2, 3 respectively and a, b, c are there opposite edge lengths, which are the same as the angles at dual vertices $1', 2', 3'$ respectively.

We can express these identities equivalently as a single one, denoting n' by n^{-1} , as follows.

$$\sum_{\epsilon_1, \epsilon_2, \epsilon_3 = \pm 1} \epsilon_1 \epsilon_2 \epsilon_3 \triangle(1^{\epsilon_1}, 2^{\epsilon_2}, 3^{\epsilon_3}) = 0$$

We can generalize these identities to a convex polygon as follows:

$$\text{vol (I)} + \text{vol (III)} = \text{vol (II)} + \text{vol (IV)} = 2\pi$$

or equivalently,

$$\text{vol (I)} - \text{vol (II)} + \text{vol (III)} - \text{vol (IV)} = 0,$$

where the regions I to IV are depicted in Fig 5.10.

Let's generalize these observations to the n -dimensional case. First start with a convex polyhedron P and its dual P^* on $\mathbb{S}^n$. Let P be the region I and consider the the region II consisting of polyhedra having $(n - 1)$ -dimensional face in common with I = P , and the region III of polyhedra having $(n - 1)$ -dimensional face in common with II, $\dots$, and so on, so that we obtain P^* as the $(n + 2)$ -th region. Then we can expect the following identity as above.

$$\text{vol (I)} - \text{vol (II)} + \text{vol (III)} - \dots + (-1)^{n+1} \text{vol (P}^*) = 0.$$

When $n = 3$, this becomes

$$\text{vol (I)} - \text{vol (II)} + \text{vol (III)} - \text{vol (IV)} + \text{vol (V)} = 0,$$

or equivalently,




Fig 5.10

$$\begin{aligned} \text{vol}(P) + \text{vol}(\text{III}) + \text{vol}(P^*) &= \pi^2 \\ \text{vol}(\text{II}) + \text{vol}(\text{IV}) &= \pi^2. \end{aligned} \quad (13)$$

Indeed the identity (13) is known as Santaló-Milnor relation, and is proved by Santaló using integral geometry for a 3-simplex [15] and proved by Milnor using Schläfli formula for a general convex polyhedron [10]. This also can be proved using elementary geometry [3].

We can obtain a simplex result for a hyperbolic convex polyhedron on $\mathbb{H}^n$ using the hyperbolic sphere $\mathbb{S}_H^n$. Consider a hyperbolic triangle $\triangle(1, 2, 3)$ and its dual $\triangle(1', 2', 3')$ on $\mathbb{S}_H^2$. Then as for the spherical case, we obtain

$$\sum_{\epsilon_1, \epsilon_2, \epsilon_3 = \pm 1} \epsilon_1 \epsilon_2 \epsilon_3 \triangle(1^{\epsilon_1}, 2^{\epsilon_2}, 3^{\epsilon_3}) = 0, \quad (14)$$

where $1^{-1} = 1', \dots$ etc. Here the dual triangle $\triangle(1', 2', 3')$ is not a convex hull of its vertices, but rather its complement to obtain a decomposition of $\mathbb{S}_H^2$ into 8 triangles as before. Also we consider only a *compact* hyperbolic triangle $\triangle(1, 2, 3)$ in the formula (14). In fact, (14) does not hold for a more general type triangle $\triangle(1, 2, 3)$, even for an ideal triangle. In these cases we have to correct (14) by some constants.

For a compact hyperbolic convex polygon, the following identities are obtained easily as for the spherical case.

$$\text{vol}(\text{I}) + \text{vol}(\text{III}) = \text{vol}(\text{II}) + \text{vol}(\text{IV}) = \frac{\text{vol}(\mathbb{S}_H^2)}{2}$$

or equivalently,

$$\text{vol}(\text{I}) - \text{vol}(\text{II}) + \text{vol}(\text{III}) - \text{vol}(\text{IV}) = 0.$$

See Fig 5.10 for the description of regions.

Again for n -dimensional case, we can expect the following identity for a hyperbolic convex polyhedron P and its dual P^* , as the complement of the convex hull of its vertices in $\mathbb{S}_H^n$.

$$\text{vol}(P) - \text{vol}(\text{II}) + \text{vol}(\text{III}) - \cdots + (-1)^{n+1} \text{vol}(P^*) = 0. \quad (15)$$

When $n = 3$, we have

$$\text{vol}(\text{I}) - \text{vol}(\text{II}) + \text{vol}(\text{III}) - \text{vol}(\text{IV}) + \text{vol}(\text{V}) = 0$$

or equivalently,

$$\begin{aligned} \text{vol}(P) + \text{vol}(\text{III}) + \text{vol}(P^*) &= \frac{\text{vol}(\mathbb{S}_H^3)}{2} \\ \text{vol}(\text{II}) + \text{vol}(\text{IV}) &= \frac{\text{vol}(\mathbb{S}_H^3)}{2}. \end{aligned} \quad (16)$$

The identity (16) can be rewritten as

$$\text{vol}(P) + \text{vol}(\text{III}) + \text{vol}(P') = 0, \quad (17)$$

where $P' = P^* \cap K^3$.

If P has an ideal vertex, $\text{vol}(\text{III})$ and $\text{vol}(P')$ become $-\infty$ and ∞ respectively. The formula (17) is proved by Santaló using integral geometry for a compact hyperbolic 3-simplex [15], proved by Suárez-Peiró using Schläfli formula for a compact hyperbolic simplex in general dimension [17], and by J. Murakami and Ushijima using a volume formula for a hyperbolic simplex [11]. Again this can also be proved by elementary geometry on $\mathbb{S}_H^3$ using Theorem 5.1 [3].


The term $\text{vol}(\text{III})$ in (17) is the sum of the products of each edge length and its dihedral angle and interpreted as mean curvature, but in the setting of extended model $\mathbb{S}_H^3$, it simply represents the volume of tetrahedra in between P and P' lying across the ideal boundary $\partial\mathbb{H}^3$. Similarly the terms $\text{vol}(\text{II})$, $\text{vol}(\text{IV})$, $\dots$, so called integral mean curvatures, are interpreted as the volumes of collections of tetrahedra sitting in between P and P' successively in the extended model $\mathbb{S}_H^n$.

APPENDIX

Let L be a "wedge region" contained in the Lorentzian part $\mathbb{L}^n$ determined by two hyperplanes $\mathcal{P}_1$ and $\mathcal{P}_2$ such that $\mathcal{P}_1 \cap \mathcal{P}_2$ is tangent to $\partial\mathbb{H}^n$ (see the picture below). The volume of L is finite if none of $\mathcal{P}_i, i = 1, 2$, is tangent to $\partial\mathbb{H}^n$, and is infinite otherwise. When $n = 2$, we already know this. Or it follows from direct calculation similar the following integrals.

Proof When $n \geq 3$, it suffices to show that the integral

$$I = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \int_1^{\infty} \int_0^{k(x_{n-1}-1)} \frac{dx_n dx_{n-1} dx_{n-2} \cdots dx_1}{(x_1^2 + \cdots + x_n^2 - 1)^{\frac{n+1}{2}}}$$



is finite for all $k > 0$ and is infinite for $k = \infty$.

We will denote $\int f \sim \int g$ when $\int f < \infty$ iff $\int g < \infty$. Consider a spherical coordinate $(\rho, \theta_1, \dots, \theta_{n-1})$ centered at the tangent point $(0, \dots, 0, 1, 0)$:

$$\begin{aligned} x_1 &= \rho \cos \theta_1 \\ x_2 &= \rho \sin \theta_1 \cos \theta_2 \\ &\vdots \\ x_{n-1} &= \rho \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \cos \theta_{n-1} \\ x_n &= \rho \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \sin \theta_{n-1}, \end{aligned}$$

where $\rho = |x|$, $\theta_i = \angle(e_i, x_i e_i + \cdots + x_n e_n)$ with $0 \leq \theta_i \leq \pi$ if $i < n-1$, and θ_{n-1} is the polar angle from e_{n-1} to $x_{n-1} e_{n-1} + x_n e_n$ with $0 \leq \theta_i \leq 2\pi$ (see [14, p.45]). Using the spherical coordinate with $\alpha = \tan^{-1} k$, $0 < \alpha < \frac{\pi}{2}$, the integral I becomes as follows.

$$\begin{aligned} I &= \int_0^\alpha \int_0^\pi \cdots \int_0^\pi \int_0^\infty \frac{\rho^{\frac{n-3}{2}} \sin^{n-2} \theta_1 \sin^{n-3} \theta_2 \cdots \sin \theta_{n-2}}{(\rho + 2 \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \cos \theta_{n-1})^{\frac{n+1}{2}}} d\rho d\theta_1 \cdots d\theta_{n-1} \\ &\sim \int_0^\alpha \int_0^\pi \cdots \int_0^\pi \int_0^\delta \frac{\rho^{\frac{n-3}{2}} \sin^{n-2} \theta_1 \sin^{n-3} \theta_2 \cdots \sin \theta_{n-2}}{(\rho + 2 \sin \theta_1 \sin \theta_2 \cdots \sin \theta_{n-2} \cos \theta_{n-1})^{\frac{n+1}{2}}} d\rho d\theta_1 \cdots d\theta_{n-1} =: I', \end{aligned}$$

where $\delta > 0$ and notice that the region $\rho \geq \delta$ being a relatively compact region inside $\mathbb{L}^n$ has obviously finite volume. Integrate with respect to ρ using the formula

$$\int \frac{x^n}{(x+a)^{n+2}} dx = \frac{1}{(n+1)a} \frac{x^{n+1}}{(x+a)^{n+1}},$$

we have

$$\begin{aligned}
I' &= \int_0^\alpha \int_0^\pi \cdots \int_0^\pi \frac{\sin^{n-2} \theta_1 \sin^{n-3} \theta_2 \cdots \sin \theta_{n-2}}{(n-1) \sin \theta_1 \cdots \sin \theta_{n-2} \cos \theta_{n-1}} \frac{\delta^{\frac{n-1}{2}} d\theta_1 \cdots d\theta_{n-1}}{(\delta + 2 \sin \theta_1 \cdots \sin \theta_{n-2} \cos \theta_{n-1})^{\frac{n-1}{2}}} \\
&\sim \int_0^\alpha \int_0^\pi \cdots \int_0^\pi \frac{\sin^{n-3} \theta_1 \sin^{n-4} \theta_2 \cdots \sin \theta_{n-3}}{\cos \theta_{n-1}} d\theta_1 \cdots d\theta_{n-1} \\
&\sim \int_0^\alpha \sec \theta_{n-1} d\theta_{n-1}.
\end{aligned}$$

Now this final integral is clearly finite if $\alpha < \frac{\pi}{2}$, and is infinite if $\alpha = \frac{\pi}{2}$, i.e., $k = \infty$. $\square$

References

- [1] Alekseevskij, D.V., Vinberg, E.B. and Solodovnikov, A.S., *Geometry of space of constant curvature*, Encycl. Math. Sc., Geometry II, Springer - Verlag, Berlin Heidelberg New York, 1993.
- [2] Böhm, B. and Im Hof, H. C., *Flächeninhalt Verallgemeinerter Hyperbolischer Dreiecke*, Geometriae Dedicata, **42** (1992), 223 - 233.
- [3] Cho, Yunhi, *An elementary proof of Santaló-Milnor-Suárez-Peiró relation for spherical and hyperbolic polyhedra*, (will be appeared).
- [4] Cho, Yunhi, *Trigonometry in extended hyperbolic space and extended Lorentz space*, (will be appeared).
- [5] Cho, Yunhi and Kim, Hyuk, *Volume of $C^{1,\alpha}$ -boundary domain in extended hyperbolic space*, (will be appeared).
- [6] Dzan, J.J., *Gauss-Bonnet formula for general Lorentzian surfaces*, Geometriae Dedicata, **15** (1984), 215 - 231.
- [7] Jo, Kyeonghee and Kim, Hyuk, *Invariant measure and the Euler characteristic of projectively flat manifolds*, J. Korean Math. Soc., **40** (2003), No. 1, 109 - 128.
- [8] Kellerhals, R., *On the volume of hyperbolic polyhedra*, Math. Ann., **285** (1989), 541-569.
- [9] Kulkarni, R. S., *Proper actions and pseudo-Riemannian space forms*, Advances in Math., **40** (1981), 10 - 51.
- [10] Milnor, J., *The Schläfli differential equality*, Collected papers, Vol. 1, Publish or Perish, 1994.
- [11] Murakami, Jun and Ushijima, Akira, *A volume formula for hyperbolic tetrahedra in terms of edge lengths*, arXiv:math.MG/0402087.

- [12] Murakami, Jun and Yano, Masakazu, *On the volume of a hyperbolic and spherical tetrahedron*, preprint at <http://faculty.web.waseda.ac.jp/murakami>.
- [13] O'Neill, B., *Semi-Riemannian geometry*, Academic Press, New York London Paris, 1983.
- [14] Ratcliffe, J.G., *Foundations of hyperbolic manifolds*, Springer - Verlag, New York Berlin Heidelberg, Graduate Texts in Mathematics (149), 1994.
- [15] Santaló, L., *Integral geometry and geometric probability*, Encyclopedia of Mathematics and its Applications, Vol. 1, Addison-Wesley, 1976.
- [16] Schlenker, J.M., *Métriques sur les polyèdres hyperboliques convexes*, J. Diff. Geom., **48** (1998), 323 - 405.
- [17] Eva Suárez-Peiró, *A Schläfli differential formula for simplices in semi-Riemannian hyperquadrics, Gauss-Bonnet formulas for simplices in the de Sitter sphere and the dual volume of a hyperbolic simplex*, Pacific J. Math., **194** (2000), 229 - 255.
- [18] Thurston, W.P., *Three-Dimensional Geometry and Topology*, Princeton University Press, Princeton New Jersey, 1997.
- [19] Ushijima, Akira, *A volume formula for generalized hyperbolic tetrahedra*, arXiv:math.GT/0309216.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SEOUL, SEOUL 130-743, KOREA
 DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-742,
 KOREA

E-MAIL: YHCHO@UOS.AC.KR
 HYUUKKIM@SNU.AC.KR